

SELECTED ABSTRACTS FROM THE 2003 NATIONAL SPELEOLOGICAL SOCIETY CONVENTION IN PORTERVILLE, CALIFORNIA¹

BIOLOGY

MOLECULAR PHYLOGENETICS OF CAVE CRAYFISH IN THE GENUS *ORCONECTES*
J. Buhay, Brigham Young Univ., Dept. of Integrative Biology, Provo, UT 84602; J. Fetzner, Jr., Carnegie Museum of Natural History, Pittsburgh, PA 15213; K. Crandall, BYU, Provo, UT 84602

The freshwater crayfish genus *Orconectes* comprises ~90 species, including 6 troglobitic species. The cave-adapted species of *Orconectes* are found along the Cumberland Plateau in Alabama, Kentucky, and Tennessee, in the Interior Lowlands of Kentucky and Indiana, and in the Missouri Ozarks. Phylogenetic examination of 75 species of the genus *Orconectes* shows that cave-adapted members are more closely related to the burrowing genus *Cambarus* than to other stream-dwelling species of *Orconectes*. Additionally, it appears that multiple cave invasions by different surface species of *Cambarus* gave rise to the various cave-adapted species of *Orconectes*. One particular *Cambarus* species, *C. tenebrosus*, is a troglophile that occurs in caves with troglobitic *Orconectes*. *C. tenebrosus* appears to be one of the ancestors to the cave-adapted *Orconectes*. *C. tenebrosus*, which can be found in caves, springs, or stream tributaries, is likely a species complex across its range from Indiana south to Alabama. Our findings support previous hypotheses on the evolution of cave *Orconectes* by a series of separate cave invasions by surface species.

CALIFORNIA CAVE BIOGEOGRAPHY

W.R. Elliott, D.C. Rudolph, D. Ubick, T. Briggs, & J.R. Reddell, Missouri Dept. of Conservation, Resource Science Division, 2901 W. Truman Blvd., Jefferson City, MO 65109

Cave life is found in carbonate rocks in the coastal ranges, Mojave Desert, Sierra Nevada, Mother Lode region, Marble Mountains area, and the lava beds of northern California. No troglobites are known from sea caves, but they contain interesting marine life. The Mother Lode region, including Tuolumne and Calaveras counties, is comparatively rich in cave-adapted species on a national scale, including one species of mite, 5 *Banksula* harvestmen, 8 pseudoscorpions, 3 *Stygobromus* amphipods, 2 millipedes, and 4 springtails. Blind carabid beetles have been studied in other parts of the state. California caves are relatively poor in Rhabdiphoridae (camel crickets) and other elements common in other karst regions. Many caves have lost their bat colonies (usually the western big-eared bat) and are threatened by land development.

MISSOURI CAVE BIOGEOGRAPHY AND BIODIVERSITY

W.R. Elliott, Missouri Dept. of Conservation, Resource Science Division, 2901 W. Truman Blvd., Jefferson City, MO 65109

The Missouri Biospeleological Database (MBD) was developed to track and analyze Missouri's rich cave fauna. Six partners pool their data, and we currently have 935 species, 1232 localities (caves and springs), and 8640 locality records. About 973 (17%) of 5800 caves have biological records, but only 408 caves (7%) meet minimum criteria for biological evaluation. There are 68 described troglobites, including 32 aquatic species or "stygobites", 15 phreatobites, and 21 terrestrial troglobites. Another 16 troglobites are undescribed or possibly are troglophiles. We know of 147 troglophiles (14 aquatic), 217 troglonexes (20 aquatic), and 478 species of uncertain status (41 aquatic).

Karst biogeographic regions include the Springfield and Salem plateaus, Ozark River Basins, Perryville Karst, Hannibal Karst, and isolated areas, such as Caney Mountain. Caves east of the Mississippi River often contain a high

er diversity of terrestrial troglobites. Troglobites are known from 367 Missouri sites, but 18 troglobites occur at single sites only. Seven possible new troglobites are from single sites. Missouri shares 46 troglobites with other states.

The "Site endemism" (SE) metric and other metrics were used for ranking and comparing caves for conservation planning. Many species and biologically important caves are being added to the Missouri Natural Heritage Database, and I am now able to produce species range maps in Arcview.

A PRELIMINARY POPULATION STUDY OF THE CAVE CRICKET, *HADENOECUS CUMBERLANDICUS* HUBBELL & NORTON, FROM A CAVE IN CARTER COUNTY, KENTUCKY
H.H. Hobbs, III & R. Lawyer, Dept. of Biology, Wittenberg Univ., Wittenberg, OH 45501

The distribution and movement of adult female cave crickets, *Hadenoecus cumberlandicus*, were evaluated in Coon-in-the-Crack Cave I (THC= 212 m) in Carter Caves State Resort Park, Carter County, Kentucky. Crickets of this parthenogenetic population were studied from February to April 2003. Crickets were individually tagged with grid-style patterns using either coral pink or blue phosphorescent paint. Hind femur length was recorded for each individual unless unavailable. After a short "paint-drying period" crickets were released to within 5 m of where originally observed.

Throughout the course of the study, 2378 *H. cumberlandicus* were processed, 1277 of which were observed only (mostly juveniles). Of the 769 tagged, 332 were recaptured, thus the recapture rate was just over 43%. Only 6 crickets were recaptured more than once (2x). Based on movement data of tagged individuals, crickets were fairly mobile within the cave, demonstrating a mean travel distance of 41 m although the greatest number of crickets moved 10-15 m. The Schumacher-Eschmeyer estimate of the population size during the study period was 5508 individuals. Crickets were most dense near areas of ceiling crevices, enlarged fractures, at intersecting passages, or in "bells." The average femur length for this population of *H. cumberlandicus* was 1.89 cm (SD ± 0.23), with a range of 0.9-2.4 cm. Assuming femur length is related to age, the population appears to consist of 3 cohorts. With nearly half the population juveniles, it is likely that the population density is continuing to increase. Numerous crickets autotomized one or both of their hind legs when handled, thus imposing potentially significant restrictions on their movement patterns and distances displaced. Loss of a hind leg(s) likely influenced their ability to exit the cave and to forage in the forest litter. While a night examination of epigeal habitats directly above the cave passages was conducted on 17 April, no crickets were observed. This, perhaps, was due to the cool temperatures of the season.

GENETIC STRUCTURE AND HISTORICAL BIOGEOGRAPHY OF STYGOBITE POPULATIONS AS THEY RELATE TO DEVELOPMENT OF HYDROLOGIC SYSTEMS: A CASE STUDY IN THE EDWARDS AQUIFER OF TEXAS & NORTHERN MEXICO
J.K. Krejca, Univ. of Texas, Integrative Biology, Austin, TX 78712

Gene genealogies are increasingly used to examine patterns of relatedness of populations of stygobites from different cave systems. Recent studies indicate that hydrology, rather than morphology, is dictating these patterns in a variety of organisms examined, including isopods, amphipods and fish. However, hydrology is typically examined only secondarily to explain relatedness, and virtually no studies exist that test the congruence of an *a priori* hydrologic hypothesis with a biological hypothesis. In addition, due to the recent abundance of genetic data and phylogenetic trees, there are newly available techniques to test tree congruence in a biogeographic framework (taking vicariance, dispersal, extinction, and other area-organism phenomena into

¹ The *Journal of Cave and Karst Studies* was unable to publish many of this year's abstracts because the files we received from the Convention's Sessions Committee were incomplete or otherwise unusable. In some cases, we were able to obtain the original abstracts and publish them, but it is quite likely that some otherwise important and qualified abstracts were not published this year or mistakes were included in some published abstracts. We regret the inconvenience this may cause some presenters. (The standards for publication of abstracts were published in v. 59, n. 1, p. 61 issue of the *Journal*.)

account). I am using these tree-matching techniques to test the congruence of a hydrological hypothesis (development of the Edwards Aquifer) with a biological hypothesis (gene trees of 2 aquatic isopods, Cirolanidae: *Cirolanides* and Asellidae: *Lirceolus*). Though analyses are incomplete, the general pattern shows congruence for recent nodes (closely related cave systems) and lack of congruence for historic nodes (cave systems in different segments of the aquifer). A likely explanation for this pattern is that the hydrologic trees are not completely accurate in their representation of the hydrogeologic development of the aquifer. This method shows great promise for studies of modern hydrologic relations, particularly in situations where traditional techniques such as dye tracing fail, as well as historic hydrologic relations that are less studied but are relevant for conservation considerations, such as contaminant transport.

CAN THE BLIND SEE? STUDIES OF VISUAL PIGMENTS IN STYGOBITIC CRUSTACEANS
M.L. Porter, Dept. of Microbiology, Brigham Young Univ., Provo, UT 84602;
A. Perez Gonzalez, Departamento de Invertebrados Universidade Federal, Rio de Janeiro, BRAZIL; M. Perez-Losada, Brigham Young Univ., Provo, UT 84602

One of the most commonly cited examples of troglomorphy is eye loss. However, previous studies of cave-adapted crayfish indicate that while the structural components of the eye degenerate, the visual pigments responsible for light capture remain functional. To determine if this observed functionality is unique to crayfish or a widespread phenomenon, the evolution of the visual pigment gene (opsin) was investigated in several additional hypogean crustacean species, including the shrimp *Antromysis cubanica* (Mysidae, Mysidacea) and the freshwater anomuran crabs *Aegla cavernicola* and *A. leptochela* (Aeglidae, Decapoda).

The *Aegla cavernicola* exhibit differing degrees of eye degeneration and the most closely related surface species have been determined as *A. strinati* and *A. marginata*, respectively. Sequenced opsin genes from both cave adapted shrimp and crabs do not show any evidence of loss of function (i.e., no stop codon accumulation or significant sequence differences).

In comparisons of the opsin gene between surface and cave pairs of *Aegla*, the opsin sequences of *A. cavernicola* and *A. strinati* were identical. In *A. leptochela*, 2 forms of the opsin gene were detected, with the calculated divergences from *A. marginati* as 0.14% and 9.7%. Although the closest surface relative has not yet been determined, the divergence of *A. cubanica* from the most closely related crustacean opsin is 2.1%.

These results indicate that in cave adapted crustaceans, the conservation of function in the opsin gene is widespread. Further studies will be aimed at determining when and where the opsin gene is expressed in crustaceans.

THE BIOGEOGRAPHY OF THE SUBTERRANEAN INVERTEBRATE FAUNA OF WEST VIRGINIA
K. Schneider & D.C. Culver, American Univ., 4400 Massachusetts Avenue NW, Washington, DC 20016

Species richness in a series of adjoining caves in Greenbrier County, West Virginia, was investigated in order to i) estimate biodiversity for this area, ii) determine whether cave size is related to the number of species, and iii) determine how many caves need to be sampled to achieve an accurate estimate of biodiversity. Eighteen subterranean invertebrate species were collected from 65 caves within a 25 km² area. Cave length was correlated with number of species. Few caves were needed to collect many of the species; 7 caves were needed to collect 95% of the species. By sampling only the largest 7 caves, 89% of the species were captured. However, the species accumulation curve did not reach an asymptote and estimations based on species rarity show that half of the species were not collected at all. Therefore, the patterns observed are approached with caution and more data and sampling are needed.

OXYGEN CONSUMPTION/BODY MASS RELATIONSHIPS OF *GAMMARUS ACHERONDYTES* & *G. TROGLOPHILUS* (AMPHIPODA) IN AN ILLINOIS CAVE
S.J. Taylor, Illinois Natural History Survey, 607 East Peabody Dr, Champaign IL 61820-6970; F.M. Wilhelm, M.P. Venarsky, & G.L. Adams, Dept. of Zoology, Southern Illinois Univ., Carbondale, IL 62901

Cave stream faunas typically occur in low densities, with species adapted to low levels of available nutrients. Contamination of cave streams with increased nutrients from surface runoff allows stygophilic species to become more dominant in the stream community, potentially displacing stygobitic

species that have lower metabolic rates. In Illinois Caverns, as in several other caves in Illinois' Salem Plateau, *Gammarus troglophilus*, a stygophilic amphipod, co-occurs with the federally endangered *G. acherondytes*, a stygobite. It has been suggested that *G. troglophilus* is displacing or out-competing *G. acherondytes* as a result of nutrient enrichment. To test the hypothesis that *G. troglophilus* has a higher metabolic rate than *G. acherondytes*, we measured the seasonal basal metabolic rates of both species in the laboratory. Slopes of the rate of oxygen consumption versus amphipod body mass differed. The relationship for *G. troglophilus* was steeper than for *G. acherondytes*, indicating that the larger *G. troglophilus* have a higher mass-specific respiration rate than *G. acherondytes*. *Gammarus troglophilus* may have a further competitive advantage because of their larger adult body size, which may facilitate greater reproductive capacity than for *G. acherondytes*. Reversing the current trend of habitat degradation will require a concerted effort on the surface to mitigate land use practices responsible for degradation of cave stream water quality.

CONSERVATION AND MANAGEMENT

BAT COUNTS AND THE FEDERAL DATA QUALITY ACT: *PLECOTUS TOWNSENDII* MATERNITY COLONIES IN AND NEAR MOWICH CAVE, OREGON 1983-2002
W.R. Halliday, 6530 Cornwall Court, Nashville, TN 37205

The Federal Data Quality Act was enacted in 2000 to ensure quality of data in federal actions. Bat counts and estimates of maternity colonies of *P. townsendii* at Mowich Cave, Oregon (obtained through 2 FOIA requests) are analyzed in the context of this law. The initial US Forest Service report form for the observation in "Summer 1983" specified an estimate of "40 bats (20 young?)". The subsequently documented birth rate for this bat at Mowich Cave is just under 100% per female per year. Thus, the seasonal 1983 population evidently was about 21 females and 19 young. However, the cited figure and date soon began to diverge from the original to as much as "50+ bats (7-2-84)". Such mis-statements reversed the projected population trend and provided inappropriate support for the controversial gating of this cave in 1992.

A somewhat similar misconstrual was published after 1999 exit counts: Those for July 26 and August 18, respectively, were 23 and 45 "big bats"; the latter clearly reflected 22 newly volants, as expected in August. A report distributed by the Umpqua National Forest (backdated to June 1999), however, attributed this August 1999 increase to re-closure of the gate's inset portal in July 1999. The 2001 season counts were less than half the 1983 baseline, the 2002 data strongly suggest that most of the maternity colony temporarily moved elsewhere to avoid the gate, and no data at all have been supplied for the 2000 maternity season. The Mowich Cave experience indicates that other statements about bat count data for Pacific Northwest caves should be revisited.

THE NATIONAL CAVE AND KARST RESEARCH INSTITUTE: DEFINING ITS ROLE IN CAVE AND KARST MANAGEMENT
L.D. Hose, National Cave & Karst Research Inst., 1400 University Drive, Carlsbad, NM 88220 ; P.J. Boston, New Mexico Inst. of Mines & Technology, Socorro, NM 87801

The National Cave and Karst Research Institute's enabling legislation directs it to pursue 6 objectives: 1) further the science of speleology; 2) centralize & standardize speleological information; 3) foster interdisciplinary cooperation in cave and karst research programs; 4) promote public education; 5) promote national and international cooperation in protecting the environment for the benefit of cave & karst landforms; 6) promote & develop environmentally sound & sustainable resource management practices.

All of these mandates relate to the management of caves, but lack a strategy for implementation. Developing such a strategy in consultation with the broad cave and karst community currently constitutes a major focus for the Institute.

While housed within the National Park Service during its current "establishment" phase, the Institute is growing into a broad coalition of government and non-government partners who will address issues affecting all cave and karst resources worldwide. Cave-specific programs and positions within the federal land management agencies mainly focus on issues impacting resources within their stewardship, whereas the Institute will provide a broad umbrella for all groups and individuals working to protect and manage these fragile and challenging terrains.

CAVE AND KARST INVENTORY OF WIND CAVE NATIONAL PARK
M.J. Ohms, Wind Cave National Park, Hot Springs, SD 57747

Wind Cave National Park contains over 20 known caves and a variety of karst features including blowholes, sinkholes, shelters, and sinking streams. Wind Cave is the park's primary resource, and, historically, all inventory, survey, and cave management efforts have focused on Wind Cave itself. The other caves and karst features have received limited attention.

At 28,000 acres, the park contains large areas that have never been investigated for caves or karst features. Knowing what caves and karst features exist is vital to their protection and to overall park management. Many of the caves contain paleontological and archeological resources, and one cave is known to be a maternity roost for a sensitive bat species. While 19 of the known caves have been surveyed, none have been thoroughly inventoried. Decisions regarding wildfire management, prescribed fire, backcountry visitor usage, non-native plant control, and maintenance activities could be influenced by information on the park's cave and karst resources.

CAVE RESOURCE INVENTORIES—GOALS AND METHODS: REFLECTIONS ON VIRGIN AND LECHUGUILLA CAVES

J.M. Queen, Dept. of Geology, Pomona College, Claremont, CA 91711

Cave resource inventories are intended to help determine the relationship between resources found and their locations. It is important that inventories: 1) be easily implemented in the field; 2) provide consistent results when practiced by different specialists; 3) not promote the inclusion of wrong data; 4) provide versatile tools for present & future data analysis; 5) not require the identification of features in the field that should be identified in the lab.

Rather, cave inventories should rely on readily made observations and agreed-to visual characteristics with non-impacting field-testing techniques.

The success of any inventory scheme depends on the structure of initial questions. If we ask "where are particular things found?", we are dealing in the "R-Mode". Under each variable we record the stations at which the feature occurs. If we ask "what is found at a particular site?", we are dealing in the "Q-Mode" and different variables are recorded under each locale. Q-Mode data management should be strongly preferred because geologic, chemical, and other variables may be entered as scalar values reflecting abundance, magnitude or significance (e.g., none-extreme, 0-5), etc., and this mode is more easily field-checked.

Additionally, Q-Mode analysis allows recognition of various factors, which account for different amounts and styles of variation in the data sets. These factors may be mapped to help managers recognize areas of potential interest, to ascertain the relationship of factors to bedrock geology, hydrology, etc., and to aid in resource management.

CONSERVATION AND RESTORATION FORUM

A LANDMARK CAVE RESTORATION PROJECT IN WIND CAVE, SD
R.D. Horrocks, Wind Cave National Park, Hot Springs, SD 57747; M.A. Reece, Lava Beds National Monument, Tulelake, CA 96134

Major trail construction work in Wind Cave began in the late 1800s. Additional trail work was done in the 1930s by the CCC and then in the 1950s by the National Park Service. Impacts created by these developments were partially restored during the 1980s and 1990s by volunteers at restoration projects and camps. The first phase of a full-scale cave restoration project was conducted during the winter of 2002-2003 along the Natural Entrance Tour Route. In preparation for this project, the extent and nature of all restoration projects along the developed tour routes were documented, and digital maps were created to assist in directing restoration activities. During the 6-month project, 36 tons of trail construction debris was removed from 0.59 km of that tour route. Before and after photos document the project. All historical artifacts discovered were documented, removed from the cave, and added to the museum collection.

This project refined the cave restoration techniques used in Wind Cave and led to development of better systems to document restoration work, estimate total tonnage of debris collected from along tour routes, and defined the process for completing restoration in the remainder of the developed cave. During the project, what we believe to be the first Cultural Landscape Survey in an NPS cave was conducted by a team of specialists who evaluated the developed portions of Wind Cave. They attempted to determine what the cultural landscapes in the cave are, define the landscapes to be protected, and provide guidance on protection measures.

REDEVELOPMENT OF MUSHPOT CAVE, LAVA BEDS NATIONAL MONUMENT
M.A. Reece, Lava Beds National Monument, Tulelake, CA 96134

Mushpot Cave is a developed cave at the Visitor Center facilities of Lava Beds National Monument. Mushpot receives ~35,000 visitors/yr. The cave has an asphalt pathway with >86 separate light fixtures designed to enhance safe movement and to highlight speleothem features. Significant problems involving the management of Mushpot Cave were identified in 1987 and recommendations for possible solutions were accepted on cave lighting, cave restoration, safety, and cave interpretation.

The Mushpot redevelopment project will provide a better visitor experience by improving the lighting of lava tube features while retaining the interpretive displays already present in the cave. Visitor safety will be enhanced by adding access and emergency lighting, as well as an even, non-slip trail. The cave environment will be protected by removing the asphalt trail and replacing it with a curbed concrete trail that will allow easier removal of lint. Algae growth will be limited by more focused lighting and motion sensors that will provide on-demand lighting.

NPS cave-lighting consultant Rod Horrocks worked with the Lava Beds staff to develop a new lighting system design for Mushpot using a 12-V system similar to those at Mitchell Caverns and Timpanogos Cave. Project work began in April with the removal of the asphalt pathway by the Monument's maintenance crew. The expected completion date is early to mid August 2003.

CULTURAL RESOURCES IN CAVES SYMPOSIUM:
 IDENTIFICATION, RESEARCH, AND CONSERVATION

CULTURAL RESOURCES IN THE CAVES OF MAURITIUS: AN ARCHAEOLOGICAL & ANTHROPOLOGICAL APPROACH

A. Chowdhury, Faculty of Social Studies & Humanities, Univ. of Mauritius, Reduit Mauritius; D.A. Shank, Continental Placer, Inc, 26 Computer Drive, West Albany, NY 12205

The paper summarizes cultural use of lava and karst caves, rock shelters, and overhangs in the islands of Mauritius and Rodrigues in the Indian Ocean. It begins with the different categories studied, emphasizing those that show evidence of human use or occupation. Particular attention is paid to use of caves by slaves who fled plantation oppression between 1642 and 1835, the period of enslavement in Mauritius. The research traces use of lava caves and mountaintop rock shelters by the slaves, discussing micro-settlement patterns, defense strategies, diet, and cultural nuances inferred from artifact and ecofact remains. The archaeological record is matched with the historical record for the region or period in question. We also note recent practices including black magic and sorcery evident in some long lava tunnels. Other uses of caves, including irrigation and as water sources, general recreation, drug abuse, and cold storage are also noted.

IDENTIFICATION & LONG-TERM MANAGEMENT OF CULTURAL RESOURCES IN MAMMOTH CAVE

G. Crothers, Univ. of Kentucky, 211 Lafferty Hall, Lexington, KY 40506; R. Ward, Mammoth Cave National Park, Mammoth Cave, KY 42259; C. Swedlund, 319 E. Poplar St., Cobden, IL 62920

Mammoth Cave contains a complex archaeological record of prehistoric exploration and mining, 19th Century nitre mining, and modification for visitor access. Historic portions of the cave also have a rich biotic potential, and the National Park Service would like to enhance the cave environment for various cave-using species. The Park Service attempts to preserve cultural resources in the cave while accommodating tens of thousands of visitors, fielding scientific expeditions, and modifying the environment to promote natural biological processes. A long-term NPS/Earthwatch inventory of cultural resources in Mammoth Cave provides baseline data to monitor impact to the resources from various activities.

Preliminary insights about potential impacts from ongoing use of the cave suggest that policing visitor activity and limiting access to sensitive resources can control most direct visitor impacts. Construction of visitor amenities (trails, lights) has a much greater direct impact, but can be mitigated through archaeological investigations prior to modification, and by avoiding fragile remains. Scientific expeditions, while limited in number, potentially have greater direct impacts because they are unsupervised and scientists may not be cognizant of their own impacts.

Scientific expeditions should be limited to problem-oriented research programs with sensitivity to the full range of vulnerable resources. Structural modification to the cave (airflow, encouraging animal access) has indirect impacts on cultural resources, for which the long-term effects are unknown. Monitoring the cave environment should begin before and continue after modifications are implemented.

SPRINGHOUSES & ROOT CELLARS: HISTORIC FOLK USES OF AMERICAN CAVES
J.C. Douglas, Dept. of History, Volunteer State Community College, Gallatin, TN 37066

The historic vernacular uses of caves in the United States include obtaining water from the environment, and storing food and liquids within it. Many caves were subsistence and domestic water sources. Evidence for this activity includes the presence of springhouses, dams, small settling structures, and piping. Caves were similarly utilized as adjuncts to domestic economies to store perishable food items. Moisture-tolerant items like eggs, milk, and cheese were preserved in wetter caves and cave springs, while drier caves or passages served as root cellars for the storage of potatoes, sweet potatoes and other items. Liquids such as cider, whiskey, and wine were also stored underground. Evidence for these uses includes cleared or partitioned spaces, wooden boxes, stone and wooden platforms or shelves, ceramic and glass containers, and doors to protect the foodstuffs from theft.

These folk practices emerged by the end of the 18th Century and continued until the middle of the 20 Century, when the arrival of electricity in rural karst areas led to their gradual demise. Common in the eastern and central U.S., these uses are also known from western caves. Vernacular environmental interactions are seldom mentioned in written sources, but oral history and geographic nomenclature can yield additional insights into these activities. Often though, material culture in caves is the only extant source. The physical evidence is, thus, important and deserving of preservation. Although little studied and underappreciated, folk uses of American caves demonstrate flexible and persistent utilitarian strains in the environmental history of caves.

CULTURAL RESOURCES IN VIRGINIA CAVES

D.A. Hubbard, Jr., Virginia Speleological Survey, 40 Woodlake Dr., Charlottesville, VA 22901

The cultural resources of Virginia caves include both prehistoric and historic contexts. Important clues to prehistoric use are torch materials and stoke patterns. Distinctive prehistoric lighting materials include cane in the more southerly latitudes, and hickory bark. Prehistoric dark-zone cultural contexts include burial, ceremonial (mud glyph and pictograph), exploration, hunting (hibernating bear), lithic, and mineral extraction. The destruction of Native American cave burial sites continues from a historic pastime to what we now know as a cultural crime. Some caves were used as long-term shelters: For example, a Russell County cave contains a 10,000-yr record of use as a habitation from the Early Archaic to Late Woodland Period.

Historic cultural contexts include evidence of cave use for archival records and self expression (inscriptions on walls), commercialization, dancing and making music, food storage (dairy and root cellar type), mineral extraction and mining (clay, onyx, & saltpetre), moonshine production, paleontological excavation (Edward Drinker Cope's workings in 1867), and water resources. Anecdotal uses include homicide evidence disposal, munitions storage (Civil War), dark-zone shelter (hiding from marauding Native Americans), and weapons disposal (Civil War).

In Virginia, prehistoric and historic cultural resources are protected from destruction or theft by the Cave Protection Act. As cavers we can best serve these cultural resources by reporting observations to appropriate researchers and cultural crimes to law enforcement authorities. Without our stewardship, the future is bleak for these resources.

PREHISTORY? WHAT PREHISTORY? ROCK CLIMBING, ROCK ART, & THE ROAD 18 CAVES

L. King, Central Oregon Cave Conservation Task Force, P.O. Box 14882, Portland, OR 97293

Between 1993 and 2002, extensive cultural resource documentation was undertaken at several Central Oregon lava tube caves developed for use as rock climbing recreation areas by local climbers. User surveys, cave mapping projects, and computer enhancement of faint prehistoric pictographs were suc-

cessfully utilized to obtain enforcement of federal protection statutes at these archaeological sites. Preservation efforts at the Road 18 Caves were actively opposed by regional climbing-related businesses and national climbing advocacy organizations. Facing a decline in available funds for archaeological site preservation and recreation management, Forest Service and Bureau of Land Management offices sought and ultimately achieved a balanced management approach in the context of problematic local recreation politics.

Documentation is merely the first step in a successful conservation project. Presentation format and timing are crucial factors in the development of cave management tools to be utilized by federal agencies.

IMAGES IN DARKNESS: AN OVERVIEW OF PREHISTORIC CAVE ART IN THE SOUTHEAST

J.F. Simek, Dept. of Anthropology, Univ. of Tennessee, Knoxville, TN 37996; A. Cressler, U.S. Geological Survey, Atlanta, GA 30319

Over the past several decades, 40 dark-zone prehistoric cave art sites have been identified in the eastern woodlands region of North America. These archaeological sites span four thousand years of prehistory and were produced by a sequence of cultures beginning with Archaic-period hunter/foragers and ending with Mississippian maize agricultural chiefdoms. We examine the environmental and cultural contexts of these sites, seeking common characteristics that allow archaeologists to predict their location and thereby manage them. Site location is considered in light of bedrock geomorphology, elevation, availability of critical resources like water, and endokarst formation and structure.

CIVIL WAR INSCRIPTIONS IN CAVES

M.O. Smith, Rt. 1, Box 461A, Rock Island, TN 38581-9627

Caves were utilized during the Civil War to an extent never before seen in the United States. Consequently, graffiti were generated in numerous caves throughout the limestone regions of the Confederacy and Border States. Some caves were used as hiding places by guerillas, robbers, refugees, Union army recruiters, and escaped prisoners of war, or as sites to secrete supplies or valuables. Many other caves were mined for saltpeter by either the Confederate Nitre Bureau or by individual contractors, and dozens of caves were toured for recreational purposes by military personnel of both sides. The study of graffiti is very important in correlating caves with specific Confederate government or contractor saltpeter mining operations. Soldier names, already found in 31 caves, may exist in nearly double that number. Civil War cave scratchings, when placed in context with traditional sources, enhance knowledge about the Confederate Mining Bureau and supplemental data given in regimental or other histories, diaries, memoirs, or official reports. Cave writings often reflect information recorded nowhere else. Therefore, they should be considered documents in their own right and worthy of preservation.

PRESERVATION OF PREHISTORIC FOOTPRINTS IN JAGUAR CAVE, TENNESSEE

P. Willey, Anthropology Dept., California State Univ., Chico, CA 95929; J. Stolen, 892 Vallombrosa Ave., Chico, CA 95926; G. Crothers, Univ. of Kentucky, Lexington, KY 40506; P.J. Watson, Washington Univ., St. Louis, MO 63130

Approximately 4500 years ago, a group of prehistoric cavers negotiated a complicated cave and discovered a passage nearly 2 hours' journey from the cave's entrance. They walked toward the back of the dead-end passage, turned around and exited the way they entered, leaving footprints and torch material behind them. These are among the earliest evidence of cave exploration in the Midlands.

Nearly 300 relatively complete footprints remained in moist mud and wet sands when the passage was rediscovered ~25 yr ago. This fragility makes the prints' preservation remarkable but vulnerable to destruction.

The modification of the prints began even as they were being made. As prehistoric cavers followed the members of their own group, they were literally walking in the footsteps of their predecessors, modifying and altering the prints as they left their own. Following the prehistoric human activities, natural processes affected the prints. When the footprints were rediscovered late last century, their destruction accelerated; modern cavers trod over the prehistoric prints. They were also altered during efforts to document the prints archaeologically, especially when a few prints were cast.

The gating of the cave several years ago offers hope that the prints will be better protected now than during the previous several decades and their

destruction slowed. By understanding the modifications, steps can be taken to protect these rare and fragile remains and we can continue to marvel at the material evidence of prehistoric cave exploration.

EXPLORATION: CALIFORNIA CAVES & CAVING

THE HISTORY OF THE EXPLORATION OF CALIFORNIA'S SHOW CAVES

A. Bosted, 2301 Sharon Road, Menlo Park, CA 94025

The discovery of gold near Sutter Creek in 1848 triggered the famous Gold Rush which brought ~100,000 hopefuls to the Golden State. Within 12 years they had panned, sluiced, dug and blasted \$550 million worth of gold from The Mother Lode area. In the process, many caves were found. Prospectors discovered California Cavern and Bower Cave in 1850, Moaning Cavern in 1851, Black Chasm in 1854, Masonic Cave in 1857, Natural Bridges in 1872, and Mercer Caverns in 1885.

The huge influx of immigrants clashed with Native American Indians, especially in Northern California, resulting in more discoveries. Kok-Chee-Shup-Chee or Natural Bridge Cave was found in 1852, Plutos was found in 1863, Lava Beds opened up in 1873, Lake Shasta Caverns in 1877, and Subway Cave around 1884.

In Southern California, Mitchell Caverns was found by silver miners in the 1860s and Boyden Cavern was found in about 1888 by loggers. These discoveries were documented by prolific Victorian writers, the most famous of which was John Muir who visited Bower, Cave City, Plutos, and Lava Beds. Most caves that are now show caves were commercialized in the 20th Century, many during The Great Depression.

EXPLORATION OF LILBURN CAVE

P. Bosted, 2301 Sharon Road, Menlo Park, CA 94025

In 1980, the Cave Research Foundation undertook a re-survey of Lilburn Cave, which at the time had a surveyed length of 12.6 km. The new survey revealed additional detail in many of the known areas, as well as entirely new sections, such as Southern Comfort, the 1 by 10 Complex, the Outback, the Opera House, Canyonlands, Area of Low Hanging Fruit, Hog Heaven, and many others. Systematic exploration was aided by the development of a detailed set of quadrangle maps of this 3-dimensional maze cave, whose surveyed length now exceeds 32 km.

TAKE IT FOR GRANITE: A SPORTING NEW GRANITE CAVE SYSTEM

Max Potter, PO Box 1425, Columbia, CA 95310

The Sierra Nevada is home to many caves formed in granite, many of them unsurveyed and largely unknown to cavers. TIFG is a high-elevation, multi-entrance cave in a drainage of the Dardanelles, a large lava-capped plateau in the Carson-Iceberg Wilderness. Piracy of a surface stream into its bed has produced a large canyon passage broken by 4 waterfalls requiring rope to descend. A lower section with a steep gradient is all climbable. Corrosional speleogens, potholes, waterfalls, and chutes characterize the passage. A survey is underway but has been hampered by the short season in which this cave can be visited.

IXL CAVE RETROSPECTIVE—FIFTY YEARS OF USE & ABUSE

B.W. Rogers, US Geological Survey MS-999, 345 Middlefield Rd, Menlo Park, CA 94025; D. Snyder, M. Haye, & I.E. Randall, Western Cave Conservancy, 213 Elm St, Santa Cruz, CA 95060

In late 1953, William Miles and caver friends of the IXL Club found cold air blowing from a grapefruit-sized opening in marble west of Santa Cruz, CA, and dug open IXL Cave. Initial exploration and excavation by the conservation-minded youths revealed ~155 m of passage, much of it crawlways. At year's end in 1966, NSS cavers digging for a second entrance revealed, instead, a pit complex dropping nearly 30 m.

Mapping has shown the cave is ~610 m long. The cave was once famous for its heavily decorated as well as size-challenging passages. However, being near a vacation-oriented seacoast town and adjacent to a University of California campus, the cave was, and still is, heavily visited by local youths as a "rite of passage". In addition, in early 1954 two newspaper articles reported the cave's discovery, and a subsequent rescue of portly "Sunday afternoon" cavers several months later unduly publicized the cave on a regional level.

As a result, the cave suffered significant damage from vandalism and carelessness. By the late 1970s, vandalism was directed at removing "magical

crystals." By the mid-1980s the cave was gutted of its formerly outstanding decorations. In 1997, the cave was obtained as an addition to Wilder Ranch California State Park and efforts at management are being undertaken. Recently, photographs taken within a few weeks of its discovery were located. These allow comparison of the former beauty of IXL Cave with its present condition. Once again, the lack of responsible ownership and lack of a management plan have essentially destroyed a once beautiful cave.

BENEATH THE DESERT: CAVES OF THE MOJAVE & THE MOJAVE CAVE PROJECT

B. Szukalski, ESRI Cave & Karst Program, 380 New York St., Redlands, CA 92373

The Mojave Desert is a vast expanse encompassing roughly 40,000 km², including much of southeastern California and portions of Nevada, Utah, and Arizona. Though not normally associated with caves and karst, California's Mojave includes major carbonate areas and some of the better-known caves in the state, including Cave of the Winding Stair, Mitchell Caverns, Kokoweef Cave, and others. The California Mojave also includes a wide variety of cave types with significant pseudokarst, granite, and lava caves. The Mojave National Preserve was established in 1994. In 2001, the Mojave Cave Survey (MCS), an informal project of the Southern California Grotto, was established to assist in the inventory and survey of caves within the preserve and surrounding areas. Recent work by the MCS has included compilation of historical data and the discovery and mapping of several limestone caves as well as numerous pseudokarst caves.

EXPLORATION: CAVE DIVING

THE QUINTANA ROO SPELEOLOGICAL SURVEY: RECENT ADVANCES IN UNDERWATER CAVE EXPLORATION IN QUINTANA ROO, MEXICO

J.G. Coke IV, P.O. Box 8663, The Woodlands, TX 77387

The Quintana Roo Speleological Survey (QRSS) supports conservation, safe exploration and survey documentation of the underwater caves in Quintana Roo, Mexico. The present study area incorporates 300 km² in eastern Quintana Roo, including 91 independent underwater caves (431 km of surveyed underwater passage). Over 100 collaborators have contributed raw survey data to the database, establishing one of the largest archives of underwater survey data in the world. A database of 230 GPS locations for underwater karst formations, and 74 km of land survey provide surface control points for an emerging regional analysis of cave relationships. Computer software converts raw cave survey data into georeferenced coordinates. Additional software manages GPS data, while calibrating topographic maps and aerial photographs.

EXPLORATION: CAVE RESCUE

A METHOD FOR TURNING A LOADED ROPE INTO A 3:1 HAUL SYSTEM

Cindy Heazlit, 5672 Bluegrass Lane, San Jose, CA 95118

The 3:1 system is considered one of the basic systems for rescue. Unfortunately, it can only be rigged when it is not under load. Most cavers do not carry an extra piece of rope for a haul system, so are in a quandary when an individual gets stuck on rope and needs to be hauled up. Use of a "long tail" progress capture device (PCD) allows the rescuer to transfer the weight of the loaded rope off the main anchor knot. The rope can then be re-rigged into a 3:1 haul system with its own PCD. The physical properties of the long tail PCD allow the load to be naturally transferred to the 3:1 PCD after the rescuers make the first haul. As this method uses minimal equipment and manpower, it has an excellent application for self rescue.

EXPLORATION: COMMUNICATIONS & ELECTRONICS

THRU-THE-EARTH 2-WAY VOICE COMMUNICATION WITH CAVE DIVERS

B.L. Pease, 567 Fire St, Oakdale, CT 06370

Commercial acoustic voice communications equipment, used by divers to talk to the surface, can be easily converted into thru-the-earth radios. For this project, 2 units manufactured by Ocean Technology Systems were converted, then used for diver-to-surface voice communications in underwater Florida caves. These 32.768 kHz USB units had ~2 Watts output. This frequency was suitable for the water-saturated limestone with its conductivity of ~.02 S/m.

The conversion involved replacing the acoustic transducers with antennas and filter/matching networks. The surface antenna was a 1-m² resonant vertical loop matched to the STX-100 surface unit. The diver's antenna was a 7.6 m "earth current" trailing wire antenna. The wire impedance was ~80 Ohms in typical Florida Springs with $s = .028 \text{ S/m}$. This was matched to the SSB-2010 diver unit, which used a full-face mask containing a microphone, earphones, and a PTT switch.

Both antennas had simple LC filters for improved RF interference rejection. This equipment was used to maintain continuous 2-way voice communication while divers explored 3 different systems at depths to 60 m. Separate 3496 Hz gear was used for tracking. At Hart Springs, divers were directed under surface features while they reported depths and their track was marked on the surface. The surface track was surveyed in 3 dimensions, in effect mapping the cave from the surface. This simple voice link has many applications.

EXPLORATION: INTERNATIONAL EXPLORATION

RECENT EXPLORATIONS IN CUEVA DEL TECOLOTE, TAMAULIPAS, MEXICO

A. Addison, 3 Sheffield Ct, St. Charles, MO 63304

Cueva del Tecolote is a major cave system located within the Proyecto Espeleológico Purificación karst study area. Explorations since 1999 have pushed the total length of the cave over 40 km. The 1999 expedition discovered significant passages in the Jellybean Junction area of the cave and a major new route between the Gargoyle Gallery and the Chihue Freeway. The 2001 Expedition added the 7 of 9 Borehole and the March 2003 expedition added almost 4 km of new survey in many areas of the cave. ArcPAD GIS software was used on a handheld computer (in cave) to manage the >100 leads for the 2003 expedition. Many leads remain to be pushed in the cave and there is a renewed interest in pushing other caves in the area.

PROYECTO CERRO RABÓN 2003

C. Andrews, 427 S. Loomis, Ste. B, Fort Collins, CO 80521

The Cerro Rabón is a limestone massif located in the southeastern margin of the Sierra Madres in the Sierra Mazateca and is well known for its deep cave potential. It is situated in the northern region of the state of Oaxaca, 300 km ESE of Mexico City, and rises 2000 m above the tropical lowlands of the Gulf of Mexico.

Systematic speleological research on a bi-annual basis began in 1985. This heavily karstified uplift of limestone has since yielded hundreds of kilometers of caves. The Kijahe Xontjoa system has been pushed to a depth of -1223 m, well below one of its speculated resurgences, the Nacimiento del Río Uruapan. An unexplored region to the south named the upper Uruapan Valley is thought to be part of the catchment area for the Río Uruapan. Its rather large high flow rate of 20m³/sec may be an indication of a large and deep cave system.

In February 2003, a team of 12 cavers from America, led by Cerro Rabón veteran Mike Frazier, ventured into the upper Uruapan Valley to search for insurgences to the Río Uruapan. One cave was mapped from previous year's reconnaissance and the team retreated with valuable knowledge to assist further expeditions into the area.

EXPLORATION IN SUPPORT OF WORLD HERITAGE STATUS FOR CAVES IN WESTERN EUROPE

K. Downey, 21 Massasoit St., Northampton, MA 01060

Several recent proposals to the UNESCO World Heritage Commission include some of the most significant caves of southern France and the karst area of the Swiss Jura. Several sites are proposed as a single group due to their unique speleothems and spectacular mineralization despite being scattered over a distance of several hundred kilometers. The Sheatenflu site is listed as a proposed international biosphere preserve. The World Heritage designation offers the cavers a powerful tool for preservation and protection of these sites, but has considerable requirements, including the mapping or re-mapping of known cave passages, exploration and documentation of entire zones of cave development, and scientific study and documentation of each site. The exploration in both some well-known and tourist caves as well as in as previously "secret" caves has been productive, and is well underway with participants from several countries. The projects are generally well supported by cavers, but are not without controversy.

THE 2003 ATACAMA HALITE KARST EXPEDITION, NORTHERN CHILE

S. Fryer & J. Despain, Sequoia National Park, Three Rivers, CA 93271; K. Downey, 21 Massasoit Street, Northampton, MA 01060; C. Walck, 10485 Courtenay Lane, Truckee, CA 96161

The Cordillera de la Sal lies at 2800 m msl in the foothills of the Andes. This 130+ km long series of ridges is composed of a series of siltstones, gypsum, and halites, which has been folded and faulted. The salt terrain has been carved into a bizarre landscape with prominent hoodoos. A brief reconnaissance trip in 2002 indicated the presence of unusual caves in the salt beds that warranted a focused effort at mapping and study. The 2003 project team included 6 cavers to focus on the exploration and mapping of caves and to determine the general potential for cave development in the salt karst. Dozens of entrances and 6 main caves were located in the halite beds near San Pedro de Atacama, an oasis town at the edge of the world's driest desert. In the Valle de la Luna area, the conditions of relatively pure halite units and rare rainfall allowed the development of impressive caves. These caves contain some spectacular secondary mineral deposits and passages range from relatively large to crawls, squeezes, and climbs. The density of caves and extent of karst indicates potential for more exploration. The caves are fairly similar to carbonate rock caves, but the walls are transparent and quite edible!

SIMA PUMACOCHA, PERU

I. McKenzie, 511 9th Street NE, Calgary, Alberta, T2E 4K4 CANADA

The lack of long or deep caves in the world's highest mountain ranges led some cavers to believe that high-elevation karsts are incapable of significant cave development. But cavers from Canada, Britain and Peru challenged this assumption in Sima Pumacocha, a newly-discovered cave located high in the Andes of central Peru. Explorations in 2001 and 2002 pushed and mapped this vertical cave down dry shafts and waterfalls to a sump at -638 m, making Sima Pumacocha the deepest cave in South America. With an entrance elevation of 4375 m msl, it may be the highest significant cave in the world.

CUEVA CHARCO: 3RD DEEPEST CAVE IN MEXICO

M. Oliphant & N. Pistole, 4105 Lowell Ave., La Crescenta, CA 91214

Cueva Charco is located in the state of Oaxaca, in southern Mexico. The cave was first looked at by cavers in 1989, and exhibited potential as a pathway into the downstream portion of Sistema Cheve. However, further exploration showed that the cave parallels the projected trend of Cheve, but drops at a steeper angle towards the groundwater level. A team of cavers set 2 underground camps in 2003 to reach the end of the cave. After kilometers of narrow, tortuous passage, the cavers encountered 300 m of borehole, and then a terminal sump. One significant side lead was pushed almost 400 m before the passage became too tight to follow. With all the data tallied, Cueva Charco is 1278 m deep and 6.71 km long.

MILLIONS OF CAVE PEARLS & OTHER CAVING ADVENTURES IN TABASCO, MEXICO

J. Pizarowicz, Wind Cave National Park, Hot Springs, SD 57747; A. Snow, Sequoia National Park, CA 93262

Further explorations in Tabasco, Mexico, have uncovered additional caves near Teapa and Arcadio Zentilla in the Sierra Madrigal, Palo Quemado on Cerro Blanco, and in the Sierra Poana. Notable among recent discoveries is Gruta de las Canicas, containing perhaps the largest concentration of cave pearls yet discovered in a single cave (estimated to be between 20-200 million cave pearls), the large passages of Solidaridad, and the high elevation caves near Palo Quemado. Mexican cavers from Villahermosa continue their explorations of Agua Blanca in the Municipio de Macuspana.

2002-2003 JOINT RESEARCH EXPEDITIONS TO THE MARIANA ISLANDS: AGUIJAN, ROTA, & TINIAN

K.W. Stafford¹, D. Taborosi², T.M. Keel¹, J.E. Mylroie¹, J.R. Mylroie¹ & J.W. Jenson³, 1) Geosciences, Mississippi State Univ., Mississippi State, MS 39762; 2) Laboratory of Geoecology, School of Environmental Earth Science, Hokkaido Univ., N-10, W-5, Sapporo, 060-080, Hokkaido JAPAN; 3) Water & Environmental Research Inst. of the Western Pacific, Univ. of Guam, UOG Station, Mangilao, 96923 GUAM

A joint research program between Mississippi State University and the Water and Environmental Research Institute of the Western Pacific conducted 3 expeditions to the Mariana Islands between June 2002 and June 2003. The

expeditions focused on the islands of Aguijan, Rota, and Tinian with individuals from Guam, Japan, Mississippi, and Texas participating. Fieldwork has yielded various cave types, including: flank margin, banana hole, fissure, pit, recharge, and discharge.

The Summer 2002 expedition (12 June 2002 - 4 July 2002) was a reconnaissance of Rota and Tinian by 7 individuals who established relationships with local governments and communities, while locating primary regions of cave development. The Winter 2002 expedition (10 December 2002 - 8 January 2003), which was primarily conducted by a 2-man team, focused on features of Tinian and Aguijan yielding 5188 m of survey. The Summer 2003 expedition (3 May 2003 - 17 June 2003) involved 11 members who continued surveys on Tinian and Aguijan and began surveys on Rota.

Logistical strategies varied with the islands investigated and the time of year: Coastal surveys were strongly controlled by sea conditions and tidal fluctuations, inland surveys relied heavily upon local guides due to dense vegetation, and access to the uninhabited island of Aguijan required the support of local government bodies. Additionally, these islands contain an extensive archeological record, including prehistoric Chamorro sites and historic WWII sites. Numerous cave and karst features show evidence of Japanese occupation during WWII, including military modification, artifacts, and occasional unexploded ordnance.

2003 SISTEMA CHEVE EXPEDITION

B. Stone, 18912 Glendower Rd., Gaithersburg, MD 20879

A 10-week expedition was fielded to Cueva Cheve (Oaxaca, Mexico) in the spring of 2003 in an effort to extend the bottom of the cave (previously -1386 m deep, 22 m underwater in the "terminal" sump). Efforts in 2003 focused on a return to the sump using rebreathers and a series of bolt climbs into the high canyon leading to the sump. A route around the underwater restriction that halted the 1991 reconnaissance dive at the Cheve sump was discovered and the sump was passed within 140 m at shallow depth. A significant stream canyon extension was discovered beyond which rapidly dropped in depth before a second sump was reached at -1472 m. A 4-person dive team ferried gear to the second sump and a 2-person recon team passed the 280-m-long underwater tunnel, surfacing in a 15-m-diameter borehole. Unfortunately, the tunnel was blocked by a breakdown collapse just 50 m further, at a point 9.3 km from the nearest entrance and a depth of -1484 m. Two km of new tunnels were discovered in 2003, including 700 m mapped in high level leads above the Wet Dreams canyon.

THE MAYAN MAZE OF ACTUN KAUA

G. Veni, George Veni & Associates, 11304 Candle Park, San Antonio, TX 78249

Actun Kaua is located under the town of Kaua in Yucatan, Mexico. In 1972, David McKenzie became the first caver to enter the cave. He found it an incredibly complex maze and, with James Reddell, mapped ~6.8 km of passage by 1975. The survey languished for over 2 decades with little new accomplished. In December 2001, January 2002, and January 2003, the project was reactivated, adding 3.8 km of survey into what may be perhaps the densest concentration of passages recorded in the world at ~1 km/100 m². It is often difficult to say where the series of closely spaced cave passages ends and broad rooms filled with bedrock pillars begin. Many cave walls are covered with primitive clay drawings. Some areas have been significantly enlarged by mining the clay for unknown purposes, probably during pre-Columbian times. Many questions remain about the cave's full extent, the reason for its radically different development compared to most other caves, including those within 3-5 km, and if the part currently known is only a branch of a far more extensive complex.

CAVES OF INDIA CONTINUE TO SURPRISE: DECEMBER 2002 US EXPEDITION TO MEGHALAYA

Michael Zawada, 10162 E. Exposition Avenue, Denver, CO 80247

During December of 2002, 10 American cavers spent 3 weeks in the northeastern state of Meghalaya, India, for the purpose of cave exploration. This was the first caving expedition organized by Americans in India. The team was working closely with several Indian cavers and one caver from the UK. Ten caves were explored in the West Khasi Hills on the Iaw-Paw Plateau near Nongnah (the Langrin area). The peculiar feature of these caves is their

formation in calcareous sandstones. The longest cave examined, Krem Maw Tynhiang, was explored and surveyed to 3167 m in length with no end in sight. Total survey of the 10 caves yielded 6600 m. The caves are also fascinating due to their unique fauna characterized by an abundance of bats (including horseshoe bat), hand-sized spiders (*Hetrapoda robusta*) and cave-dwelling snakes, tentatively identified as genus *Elaphe*. Additionally, 2 blind pits in rather pure limestone were investigated in Wahlong, East Khasi Hills (the Shella area) with depths estimated at 20 & 60 m. In conjunction with cave survey, GPS, photographic, and geologic data were collected during our visit. Certain aspects of our work are available for viewing at:

<http://www.deepcaves.net/India02/india02.htm>.

In summary, Meghalayan karst once again has proven to offer extraordinary diversity for speleological endeavors.

EXPLORATION: SURVEY & CARTOGRAPHY

UPDATE ON 3D MAPPING OF A PORTION OF LAUREL CAVERNS, PENNSYLVANIA
B. am Ende, 12801 Climbing Ivy Dr., Germantown, MD 20874; L.S. Nyland, Univ. of North Carolina, Chapel Hill, NC 27599

In June 2002 a portion of Laurel Caverns, PA, was scanned with a DeltaSphere-3000 device built by 3rdTech, Inc. in Chapel Hill, NC. It is a scanning, time-of-flight laser rangefinder that takes up to 20,000 measurements/sec. The laser is directed around the environment with mirrors and motors, capturing 360° panoramas that have a 150° vertical span at a resolution of about 13 samples per degree (in each direction). The laser measures distances to the first surface it hits, up to about 15 m away, with an accuracy of ~8 mm. To overcome occlusions or shadows, range scans are taken from multiple locations. Sufficient overlap between scans allows them to be aligned with one another during post-processing.

For each range scan, there is also a color panorama acquired with a Kodak Professional Digital Camera. A wide-angle lens is used, providing sufficient color imagery at a resolution that is close to that of the range scan. Average time for a scan is ~30'. To complete the work, we spent 6.5 hrs scanning from 9 locations.

Post-processing has involved several steps. The color images are fused together with the range data, coloring each range sample. We use the Polyworks software tools from Innovmetric coupled with some of our own custom tools.

COMPASS/INCLINOMETER USING LASER POINTER FOR CAVE SURVEYING

D. Andreatta, 6041 Woodmoor St., Columbus, OH 43229

A new device has been developed that incorporates a small laser to make compass and inclinometer use easier. The laser is used as a pointer, not as a distance measurement device. The device is a 2-piece folding unit with a laser pointer and inclinometer on the front piece and the compass on the back piece.

To use the compass, the rear half of the device is approximately leveled with the back at the "from" station, and the front half tilted so that the laser points as the "to" point. The vertical plane containing the survey points also contains the line of laser light and the axis of the compass. The compass is read in the usual manner. A telescoping extension is useful when the "from" station is on the ceiling of a passage. To read inclination, the hinge of the device is held on the "from" survey station, and the laser is pointed at the "to" survey station. The inclinometer is read in the normal manner. Two types of inclinometers were tested, an electronic version and a mechanical version. Both can be used without looking at them, and hold their readings once they are set.

The device is ~15 cm long when folded and ~5 cm in "diameter". The device has been used in cave surveys and has performed well. The device has been tested around a 4-sided loop, and backward and forward loops have closed with ~1° of average error horizontally and 0.3° vertically.

CARTOGRAPHY OF LILBURN CAVE

P. Bosted, 2301 Sharon Road, Menlo Park, CA 94025

Lilburn is a 32-km-long, 3-D maze cave that presents interesting cartography challenges. This has led to the development of a multi-level quadrangle system. Approximately 90 quadrangle maps are needed to cover the entire cave. For convenience in reproduction and in-cave use, the text and level of detail are sized for 11" by 17" format, but can also be used in 8.5" by 11" format. Up to 4 vertical slices are made for each plan view. The slices are selected on a station-by-station basis for the best clarity, rather than strictly by ele-

vation. A special code for each station is used by a custom plot program to decide which level(s) a given station is to be plotted on. Typically, 3-5 stations of overlap are used at level transitions. Each transition is given a number, and prefixed by "U" or "D" to tell the reader whether to look "up" or "down" a level for the corresponding number. A program was written on a Mac to make the line plot and borders given the code for a given quadrangle. The cave passage and details are drawn in Canvas or Illustrator. This makes it much easier to make changes than using pen and ink. Additional maps to visualize the overall cave have been made with custom software, including stereo 35-mm slide pairs.

REGIONAL STUDIES FOR UNDERWATER CAVES IN QUINTANA ROO, MEXICO
J.G. Coke IV, P.O. Box 8663, The Woodlands, TX 77387

Underwater cave survey techniques integrated with the Global Positioning System (GPS) disclose underground drainage configurations and cave relationships in eastern Quintana Roo, Mexico. The present study area incorporates 6300 km². Garmin II+ and Garmin V GPS receivers, calibrated for civilian applications, verified entrance coordinates at independent underwater cave systems. With Selective Availability disabled, a 12-channel receiver reports a 3-9 m Estimated Position Error at coordinate collection locations. Correlating distant cave systems by conventional land surveys in the indigenous jungle environment is problematic; those surveys exceeding 1.5 km encourage GPS applications. A database of 230 GPS locations for underwater karst formations, and 74 km of land survey provide surface control points for the regional analysis.

Safety and environmental issues require underwater cave explorers to amend traditional cave surveying methods. A permanent guideline, knotted at fixed intervals, is positioned in new passage during initial exploration. Survey data are collected on the exit utilizing a depth gauge and compass, while knotted segments of the guideline are tallied between survey stations. Eleven underwater caves in Quintana Roo are surveyed by more accurate methods. Taped survey lengths, compass backsights, and measuring passage attributes are means used to create these maps.

THE EFFECTS OF LAVA ON COMPASS READINGS

D. Green, 4230 Sovereign Way, Salt Lake City, UT 84124

Cavers mapping lava tubes know well that compass readings are not always what they should be. This is, many times, incorrectly attributed to the attraction of the magnetized compass needle to magnetic material (magnetite) contained in lava. However, the main causes of unwanted needle deflections are due to distortion of the magnetic field because: 1) a property of magnetic material called susceptibility and; 2) magnetization of magnetite from lightning strikes.

Knowing how these effects arise and their characteristics may aid the mapper in achieving better readings. Fore- and back-sights cannot correct for readings caused by distortion of the earth's magnetic field.

DJUVU MAP ENCODING FOR DISPLAY ON THE WEB

M. Passerby, P.O. Box 80693, Lansing, MI 48908

Using DJVU encoding has given the Raders Valley Project the ability to communicate visually to project participants using highly zoomable, crisp, small file size images of current working maps. Further information such as "zoom startup" and embedded hyperlinks in the image further the usefulness of this technology. To begin the process, cave project participants must download a small browser plug-in that will allow the embedded images to be loaded via their web browser. Then maps both working and complete can either be scanned or opened in the DJVU editor and encoded for display on the web. Using the zoom tools in the browser, the project participant can now zoom in to specific areas with very little loss of detail to focus on areas of interest within the survey. This technology opens the door to sharing maps with incredible detail and a small file size via the web.

MORPHING DIGITAL WORKING SKETCHES TO LINE/WALL PLOTS USING MANIFOLD 5.5

M. Passerby, P.O. Box 80693, Lansing, MI 48908

Registering and morphing a rough digital working sketch to a line plot only accomplishes part of the complex process needed to accurately trace a final digital cave map. To fully accomplish a digitally "traceable" bottom layer requires that the passage walls from section to section match up and further

that the left and right points at each station are registered and morphed along with the sketch to the line plot points. I have developed such a workflow process using Manifold 5.5. First, working sketches are scanned and then opened in Manifold. A separate lineplot image with tics for left/right is then opened in the same project and control points placed on both images at spots corresponding to survey stations, left/right points and wall segment joining points. Using a geo-registering method that Manifold calls "Affine", the rough sketch is then morphed onto the line plot and the assembly begins piece by piece. The final result is a fully aligned and digitally traceable cave map layer. Experience has shown that proper order in placing the control points results in consistent and highly accurate results.

EXPLORATION: US EXPLORATION

WIND CAVE: 100 MILES AND BEYOND

C. Bern, 12722 W Virginia Ave, Lakewood, CO 80228

The survey of Wind Cave (Wind Cave National Park, SD) grew from 150.8 to 171.2 km in 2001 and 2002. This moved the cave from 8th to 6th on the world's longest caves list. Achievement of the 161-km mark happened in August 2001. Exploration was accomplished by Park personnel as well as volunteers, including cavers from Colorado's Wind Cave Project. Trip leaders focused on a variety of areas in the cave. Exploration occurred in the Club Room, Colorado Grotto, Halfmile Hall, Historic, North, and Southern Comfort sections. The pace of exploration has been impressive: 12.1 and 8.3 km were mapped in 2001 and 2002, respectively.

Cavers pushed the edges of the map and these efforts met with some success. Overall, however, the known boundaries of the cave changed little. Many of the 20.4 km discovered were in the interior of the map. An example is the Kneebone area, originally discovered in 2000. This area has yielded >3.6 km of passage and filled in a significant void in the map. Numerous smaller discoveries included geologic features, historic artifacts, water inlets, and a near-connection with the surface. The keys to success have been systematic checking of leads. Many miles remain to be explored, and it is difficult to predict where they will be found.

EXPLORATION OF THE KIPUKA KANOHINA CAVE SYSTEM, HAWAII CO, HAWAII

D. Coons, 586 E. 9th Rd., Rutland, IL 61358

South Point on the Big Island of Hawaii is aptly named, because it forms the southernmost point of land in the United States. A prehistoric pahoehoe lava flow named the Kipuka Kanoehina is just a few miles north of this windswept peninsula. Exploration and survey within this geologic unit has revealed one of the longest and most complex basalt cave systems known to the world. To date, 25.52 km of survey has been accumulated. More than 40 open leads remain to be explored.

EXPLORING EASTERN KENTUCKY'S COAL TRACE CAVE

J. Despain, Sequoia National Park, Three Rivers, CA 93271

Coal Trace Cave was found by Art Petit in the late 1990s in Wayne County, KY. The cave lies above a spring that flushes coal (presumably from nearby strip mines) during large floods. Coal Trace features large passages, a prominent stream, strong airflow, and very tight breakdown chokes. An effort to survey the cave led to the discovery of the Catron Chamber (named for the landowner), which at its largest is 52 m wide and 27 m tall. Although this section of the cave was touted as the largest cave room in Kentucky in numerous newspaper articles around the country in 2000 and 2001, the area is really a large borehole. Currently the cave has been surveyed to a length of 2.4 km, several very good leads remain, and a connection to nearby Barefoot Saltpetre Cave is possible.

EXPLORATION IN CARROLL CAVE, MISSOURI

E. Frantz, 8468 N Overland Ct., Kansas City, MO 64154

Carroll Cave has been closed by the landowner for over 15 years. With >19 km of surveyed passage and at least 100 known but unsurveyed side passages, Carroll Cave is one of Missouri's most significant caves. In December 1999, an attempt to dig into the cave through a nearby sinkhole was abandoned after 5 years of work due to safety concerns. In November 2000, a 23-cm well shaft was drilled from the surface into the cave. Mostly with the help of explosives, this shaft was expanded to 91 cm., reopening the cave in July 2002.

HYDROGEOLOGIC INVESTIGATION OF THE CARROLL CAVE AREA, CAMDEN COUNTY, MISSOURI

M.D. Hall, 5226 Slashwood Ln., Spring, TX 77379

A multidisciplinary approach is currently being used to characterize the Carroll Cave System. Mapping of surface karst and geologic and hydrologic studies in Carroll is helping direct current exploration and surveying efforts. The examination of rock samples and 9 thin-sections from the new entrance borehole of Carroll places the majority of cave in the lower Ordovician Gasconade Formation. Cavern development has been strongly influenced by structural deformation of strata. Groundwater flow follows dip of geologic structure in the Upper Thunder River and Carroll River passageways, but trends along strike of geologic structure in the lower Thunder River passage. Analysis of groundwater samples taken from various locations in the cave indicate a change in dissolved mineral saturations that may help explain the cave's development. Inorganic water quality of the cave is excellent. Flow and chemical analysis of groundwater points to extensive, unentered portions of cave. Overburden and sinkhole location maps are used to infer possible boundaries and extensions of the cave system. A surface temperature survey and characterization of a sinkhole collapse has led to an interpretation of a significant extension of the cave towards Barnett Hollow ~2000 m west of Carroll passageways. Several sinkhole valleys filled with alluvial gravels, that may support recharge to Toronto Springs, have been identified. The validation of computer geologic mapping is currently ongoing with 30 sinkholes out of hundreds field-located and photographed.

EXPLORING HYPERTHERMAL CAVES IN KILAUEA CALDERA, HAWAII

W.R. Halliday, IUS Commission on Volcanic Caves, 6530 Cornwall Court, Nashville, TN 37205

Under National Park Service permit, members of the Hawaii Speleological Survey have been locating, exploring, mapping, and inventorying caves in Kilauea Caldera since 1989. Almost 200 lava rise, lava tube, hollow tumulus and other caves have been identified in the 1919 "Postal Rift" Lava Flow. Many are hyperthermal; a few even exceed the 70°C maximum of our digital read-out thermometers. Limiting factors in exploration are temperature, humidity (steam) and subsurface winds, and fumes. Fumes are a problem only in 2 very small caves adjoining a fumarole field; elsewhere the fumes have a pleasant taste and seem to be benign. Hot and cool swirls and currents of air change quickly with changes in surficial wind direction and velocity. Thus, one-minute escape routes are maintained at all times. Thermostratification commonly is marked, with crawling necessary in some caves. Different individuals have slightly different tolerances to hyperthermia in the presence of 100% relative humidity. Empirically developed guidelines for exposures up to 52°C provide a small margin of safety for young, healthy cavers. Steam temperatures vary considerably, and the mere presence of steam no longer is considered a limitation. Personal sublingual thermometers reveal whether extensive rest and replacement of fluids and electrolytes are needed after exiting.

RECENT EXPLORATION IN LECHUGUILLA CAVE

J.T.M. Lyles, PO Box 95, Los Alamos, NM 87544

Lechuguilla Cave was reopened for exploration after a new airlocked entrance was completed in mid-2001. Cavers focused on cleaning up old surveys that had loops with poor closure or did not have reverse shots to catch blunders originally. Re-examination of crawls, climbs, pits, breakdown, and boneyard holes that were passed up or missed by earlier surveyors yielded new passage. Results of recent expeditions have extended the cave length to >175 km of total passage.

DIGGING & THE CAVES OF RADERS VALLEY

M. Passerby, P.O. Box 80693, Lansing, MI 48908

A single well known blowhole was the area's major indicator that something great lay below, but until recently not much else was known about the caves, geology, and hydrology of this magnificent dry valley in West Virginia. Now through various means of digging and micro-shaving/muzzing many caves including Caves.com Cave, Zicafoose Blowhole, Deels Hole, Bobcat Blowhole, Middle Earth and Freelanders Well are in the process of uncovering and revealing the incredible beauty of some of this area. The project has become a model for using technology to communicate and visually represent data online so project members can quickly determine where to continue

exploration both above and below ground. Exploration above ground has been accelerated by the use of a GIS system called Manifold 5.5, which has been adapted to "register" rough working sketches to the actual line and wall plots and to georegister relevant DEM's, aerial photographs, and topo maps for online display.

CAVING IN THE MAMMOTH AREA OF YELLOWSTONE NATIONAL PARK, WY

J. Pizarowicz, Wind Cave National Park, Hot Springs, SD 57747; A. Snow, Sequoia National Park, CA 93262; D. Lester, 4329 E Peakview Ct., Centennial, CO 80102

Several caves, including McCartneys Cave, Jeweled Cave, Devils Kitchen and AJs Chasm were explored and mapped during an expedition to Yellowstone National Park's Mammoth Hot Springs area in November 1999. Several of the caves were very warm (50+°C) and toxic levels of CO₂ necessitated the use of SCBAs to explore beyond some of the caves' entrances. Caves were noted to have formed in laminated, hydrothermal travertine; and microbial activity was recorded in hydrothermal pools in the caves.

JEWEL CAVE EXPLORATION

M. Wiles, 1607 Gray Blvd, Newcastle, WY 82701

Jewel Cave's barometric airflow has been a valuable tool for discovering new passages. In late 1991, a strong breeze encouraged the excavation of The Stopper, leading to the discovery of nearly 48 km in just 6 years—including larger rooms and passages than any previously known. Today, this breakthrough is less than halfway to the end of the cave.

For 30 years, exploration trips had always been done in a single long day. But with trip times approaching 20 hrs and round trip distances nearing 16 km, it became necessary to establish a single permanent camp in 1997 to facilitate continued exploration of the farthest reaches. Since then, 4-day trips—limited only by the amount of urine that can be carried out—have become common.

Strong breezes led to a second breakthrough later in 1997, and it is beyond The End that most of the cave is expected to be found. The cave currently has >208 km of mapped passages, and innumerable open leads.

GEOGRAPHIC INFORMATION SYSTEMS & DIGITAL MAPPING

ARCPAD GIS MOBILE SOFTWARE IN CUEVA DEL TECOLOTE, TAMAULIPAS, MEXICO

A. Addison, 3 Sheffield Ct, St. Charles, MO 63304

The March 2003 expedition to Cueva del Tecolote cave system in the Proyecto Espeleológico Purificación karst study area made extensive use of ArcPAD GIS software. The expedition had 23 cavers in 7 survey teams to send to over 100 possible leads. The GIS project consisted of the existing cave survey as a layer imported from WALLS cave surveying software, a layer of all other known caves in the area, a scanned version of the topographic map, scanned survey notes, and a shapefile of all the leads for the expedition. Leads were symbolized as point features and linked to the corresponding survey note scans. All of the data were loaded onto an iPAQ handheld computer for the expedition. A special "wetbag" designed for use with computers was used to eliminate humidity and water damage to the unit. The iPAQ was used for six days at Camp I and also on long survey trips to different areas of the cave.

Using ArcPAD on this expedition was a big success. Teams with little to no familiarity with a given area of the cave could simply choose a lead on the handheld, and retrieve the survey notes for the area. Leads were color coded based on the type and quality of lead, allowing survey teams to be as productive as possible. Having the handheld computer in the cave also provided programs such as Word and Excel for writing trip summaries and correcting math errors while in Camp I.

ASSESSING PERENNIAL DRAINAGE DENSITY IN THE HIGHLY KARSTIFIED TURNHOLE BEND BASIN, KENTUCKY

A. Glennon, Dept. of Geography & Geology, Western Kentucky Univ., Bowling Green, KY 42101

Drainage density is the summed length of all streams in a basin divided by the basin area. It is a measure of average length of streams per unit drainage area and describes the spacing of the drainageways.

Initially, 4 methods were developed to estimate perennial drainage density within the highly karstified Turnhole Bend Basin, KY, resulting in values ranging from 0.24 km/km² to 1.13 km/km². Since mapped cave streams repre-

sent only a fraction of the total underground streams within the study area, the actual values are likely much higher. With each method, by increasing detail, higher perennial drainage density values resulted. Therefore, efforts were taken to devise a model that would reflect a reasonable upper drainage density value.

Using ArcInfo 8.3 and Visual Basic.net, an idealized karst drainage model was developed. Using a square drainage basin, a drain point along the basin's perimeter is randomly selected. A number of sinkhole drains (x) are randomly located in the basin. Repeatedly running the model, a least-distance stream network yields a pattern consistent with the formula.

Anhert & Williams (1998) counted an average of 74 sinkholes/km² for the Turnhole Bend Basin Sinkhole Plain. Applying the provisional equation, perennial drainage density is 6.83 km/km². Though this value is derived from an idealized model, quantitative measures of subterranean stream networks hold potential as powerful analytical tools in the understanding of karst.

INEXPENSIVE LINEAMENT ANALYSIS FOR DISCOVERING INGRESS NEXUS (ILADIN) *F. McDonough, 8137 Kelvin Ave, Winnetka, CA 91306*

An inexpensive technique has been developed that increases the probability of finding cave entrances in undeveloped karst and pseudokarst terrain. The first trial of this technique was done in Sequoia National Park in late 2002. A small solution cave previously unrecorded by the National Park Cave Survey was found in the area (~465 m²) ILADIN predicted would have a high probability of containing cave entrances. Three choked solution passages, one with moderate airflow emanating from it, were also found in ILADIN highlighted areas. ILADIN's second trial, in an area known to form sandstone pseudokarst caves, also yielded a series of promising cave entrances.

WALLS 2D: REALISTIC DRAWING & MORPHING OF CAVE WALLS & PASSAGE DETAILS & ITS APPLICATION TO GIS

D. McKenzie, 16200 Sweetwood Trail, Austin, TX 78737; G. Veni, George Veni & Associates, 11304 Candle Park, San Antonio, TX 78249

Walls is a Windows-based program designed to help cave surveyors organize their data and prepare maps. Such programs typically adjust the data to compensate for inevitable errors, but Walls provides tools to identify gross errors in specific survey traverses. While some programs also draw rough approximations of cave walls based on measured distances from survey stations, Walls 2D is a new component that makes Walls the first program to realistically draw cave passages and details. Scalable Vector Graphics file formats allow the use of commercial drawing programs to create fully illustrated maps that Walls can adjust as the cave survey evolves, eliminating the need to fully or partially redraw maps. Sketches are morphed in appropriate ways, depending on their type of information, to reflect changes in the surrounding survey due to corrections and least-squares adjustments.

Certain Walls files can be exported to ArcView and ArcMap for GIS analysis and merged with track-logs and waypoints from Garmin GPS receivers to create complete, detailed georeferenced datasets. The Walls 2D output maps can potentially lead to automated and accurate calculation of cave areas, cave area to bedrock ratios, and analysis of certain hydrogeologic parameters.

USING A GEODATABASE NETWORK TO DETERMINE BEST TRAVEL ROUTES THROUGH JEWEL CAVE

R. Ohms, Jewel Cave National Monument, Custer, SD 57730

The cave survey lines and stations shapefiles generated by CaveTools were imported to an ESRI ArcGIS feature dataset, and a geometric network was created in order to find best travel routes through Jewel Cave. The path taken from the elevator entrance to Camp Duh was determined to be 7.0 km, and from the elevator to the end of current survey is 10 km. Restrictions can be placed on the trace in order to avoid certain delicate speleothems or tight constrictions, and the resulting path can be exported as a new GIS layer. This could be a useful tool for cave rescue, as the most desirable path in a rescue situation is not always the shortest. The network is currently limited by the completeness of the cave survey, passage dimension data, and feature inventory, and will be improved upon as more data are collected and entered.

KARST OCCURRENCE & SINKHOLE GIS COVERAGES FOR KENTUCKY

R. Paylor¹, L. Florea², M. Caudill¹ & J. Currens¹; Kentucky Geological Survey¹ and Kentucky Speleological Survey², 228 Mining & Minerals Resources Building, Lexington, KY 40503

Growing interest and need for better karst maps for Kentucky has prompted the development of digital karst data. The Kentucky Geological Survey (through a published 30'x60' karst groundwater basin map series in cooperation with the Kentucky Division of Water) and the Kentucky Speleological Survey (through a digital karst feature database) have independently collected and archived karst data for the past several years. Present publications based upon these data and improved by current digital geologic mapping techniques at the Kentucky Geological Survey include a Karst Occurrence map of Kentucky (depicting regions with potential for karst development) and a digital sinkhole coverage for Kentucky (consisting of karst-related topographic depression polygons). These maps are an important foundation for further work and can provide an important resource for those working in karst prone areas.

GEOLOGY & GEOGRAPHY

BIT DROPS (DRILLED VOIDS) AS A MEASURE OF KARST

R.L. Bixby & I.D. Sasowsky, Dept. of Geology, Univ. of Akron, Akron, OH 44325

Karst is typically defined based upon the presence of sinkholes, caves, and conduit springs. In areas where glaciation has occurred, however, till and other deposits may bury the surface, leaving little evidence of karst. The carbonate aquifer behavior may retain characteristics of karst (conduit, or quick flow) or may be so occluded by sediments that conduit flow does not occur. In order to develop strategies to protect groundwater quality, it is important to know which type of behavior is present.

A band of Devonian and Silurian limestone underlies central Ohio. In several areas, a subtle karst is seen at the surface and rapid groundwater flow is known to occur. In an attempt to discern the presence of subsurface karst, we evaluated bit drops (voids encountered during the drilling of water wells) for 9 counties, 7 of which were considered to have established karst. The data source was the Ohio Dept. of Natural Resources, which provides on-line access to records of 725,000 water wells in the state. In the 9 county area, 75,336 wells were evaluated for voids, and then located by township. For some of the townships not having traditional karst features, significant voids were nevertheless found at depth. A ratio expressing the number of wells with discontinuities to the total number of wells found higher values for some counties with little surface karst. This suggests that rapid groundwater flow can occur in these areas, and that they should probably be considered as "karst" for purposes of groundwater quality protection.

EXTRATERRESTRIAL CAVE-FORMING MECHANISMS

P.J. Boston, Cave & Karst Studies Program, New Mexico Inst. of Mining & Technology, Socorro, NM 87801

Earth caves form in many ways ranging from solutional processes to melting/resolidification. Numerous cave types derive from a relatively limited number of geologic and physical processes identified over time by speleologists. Using this foundation, the physics of speleogenetic processes can be extrapolated to other bodies in the Solar System. This type of thought experiment is essential to bolster advocacy of caves as important scientific targets for missions to other planets and moons. Results of such informed speculation are utilizable by mission planners. It can also sensitize non-speleological scientists to possible cave occurrences on extraterrestrial objects.

Planetary mission images show visually detectable lava tubes on Mars, the Moon, Venus, and possibly Io. Other inconspicuous cave types often have no or only subtle surface indicators. Geophysical techniques can elucidate subsurface structure from orbital, aerial, or lander platforms. All methods have advantages and limitations viewed in light of particular missions.

Rocky and icy bodies can experience internal and external energy sources resulting in cracking, surface material shifting, melting, and resolidification. For example, on bodies without liquid water, other solvents (e.g., liquid ammonia, nitrogen, or ethane) may create unique speleogenesis. Subliming water/CO₂ ices on Mars may cause unique voids. Comet perihelion passage may produce vaporization caves.

Many have developed cave classifications. We use the dominant physical process: 1) solutional, 2) erosional, 3) tectonic (including crater impact), and

4) phase transition. Within these categories, we fit known speleogenetic mechanisms while accommodating possible unique extraterrestrial caves (e.g., cryovolcanic icetubes, Martian ground-ice sapping voids, and others).

GENESIS & SEDIMENTATION OF WINDY MOUTH CAVE, WEST VIRGINIA
M.D. Curry & I.D. Sasowsky, Dept. of Geology, Univ. of Akron, Akron, OH, 44325; D.A. Shank, Continental Placer, Inc., 26 Computer Drive West, Albany, NY 12205

Windy Mouth Cave is an 11-kmlong abandoned spring conduit formed within the Hillsdale and Sinks Grove Limestone adjacent to the Greenbrier River. The cave contains complex clastic (fluvial) sediments, which were sampled throughout the cave from gravel bars, floor deposits in domes, and main trunk passages. Particle size distributions revealed 2 end members: a high percentage of gravel, lacking mud and sand, or a high percentage of mud, with no gravel or sand. Four of the 16 samples were intermediate between the two. Chert dominated the samples, with the gravel size fractions averaging 75%. Quartz (as determined by XRD) dominated the clay fraction lithology. Grain shape percentage among size fractions >2.00 mm and >1.00 mm both contained high amounts of sub-rounded to sub-angular grains. Sediments found in the cave are locally derived from rocks of the Greenbrier Group and Maccrady Shale.

Sinkhole inputs from the surface feed different branches of the cave. Most sinkholes have very limited catchment in the present environment, and are underfit in relation to the paleodischarges recorded in the sediments within the conduits. Chaotic gravel deposits indicate periods of intense, probably pipefull, water flow. Fine-grained deposits relate to low flow conditions caused either by lack of recharge or obstruction of the conduit outlet. An ill-defined magnetic reversal found in sediments near the Waterfall Room suggests an age of >760,000 years for deposition of those materials. This is consistent with the height of the cave above present-day base level (30 m).

A HIDDEN SPELEOTHEM: ELONGATED CONCRETIONS IN COLORADO CAVE SANDS
D.G. Davis, 441 S. Kearney St., Denver, CO 80224; F.G. Luiszer, Dept. of Geological Sciences, Univ. of Colorado, Boulder, CO 80309

While digging out ancient sediment chokes to extend 4 partially filled caves in Colorado (Cave of the Winds, Sandite Cave, and Cave of the Swirling Mists in Williams Canyon; Lost Mascot Cave in the Fulford district), cavers have unearthed hidden deposits of elongated concretions of calcified sand (informally called "sandites" by the diggers), embedded in otherwise unconsolidated sand. These are of circular to flattened cross-section, sometimes curving gently or splitting into parallel forks, and may have either rounded or pointed ends. They are up to ~60 cm long and from 6 mm to 5 cm or more wide. Many are isolated and unattached; others are prong-like extensions of larger, irregular calcified masses. In most locations, they parallel the axis of the filled passage in a slanting to subhorizontal attitude, and may cut across bedding layers. They probably grew when the sand in filled sumps was locally saturated with calcite-rich water, but not necessarily below the regional water table. Similar growths have been reported in Australian caves, but not, to our knowledge, in other American ones. Deposits of comparable elongated concretions have also been found in non-cave sands and sandstones in the U.S. and Italy. The calcite-deposition mechanism is undetermined, but the cave examples seem to confirm one conclusion published by researchers of the non-cave concretions: The long axes tend to be aligned along the groundwater flow direction.

HYDROGEOLOGY OF SPENCER MOUNTAIN: LICK BRANCH KARST INVESTIGATION
J. Deatrick, US EPA - SESD/EAB, 980 College Station Road, Athens, GA 30605; J. Hoffelt, Tennessee Division of Superfund, 711 R.S. Gass Blvd., Nashville, TN 37216; W.S. Anderson, K2 GeoEnvironmental, Inc., P.O. Box 210485, Nashville, TN 37221

Karst hydrogeology investigations continue around Spencer Mountain in Van Buren County, Tennessee. The court-ordered settlement regarding treated effluent from the City of Spencer's wastewater treatment plant required a karst investigation of the Lick Branch to Pennywinkle Spring drainage basin. A temporary discharge of treated effluent is allowed into the Pennywinkle drainage while a pipeline is built to the Caney Fork River. The purpose of the karst study is to determine the potential for treated effluent to reach water supply wells and sensitive cave ecosystems.

The Tennessee Dept. of Environment and Conservation requested that the U.S. EPA Science and Ecosystem Support Division lead the study. Members of the National Speleological Society provided volunteer assistance, and K2 GeoEnvironmental provided specialized analytical services.

A June 2002 dye study used field and laboratory fluorometry analyses of water and passive receptors. Base flow conditions existed at the start and end of the study with 3.94 cm rainfall 12 hrs following initial dye injection. The study confirms that water in Lick Branch flows underground through Lick Branch Cave. The water resurges on the Hartselle Formation, flows over a waterfall, and sinks in the Monteagle Limestone. The water resurges again at Pennywinkle Spring in a distributary spring fashion. Sampling and analysis of area water supply wells did not confirm the presence of dye in any drinking water system. Dye mass recovery at Pennywinkle Spring was 97.5%. Another dye trace showed that Lost Cove (a karst valley) water flows through the Monteagle Limestone to Pennywinkle Spring.

CONSEQUENCES OF LOW pH, CAVE-WALL CONDENSATION & BIOFILM DEVELOPMENT TO SULFURIC ACID SPELEOGENESIS

A.S. Engel, L.A. Stern & P.C. Bennett, School for Geosciences, Univ. of Texas, Austin, TX 78712

Volatilization and oxidation of H₂S to sulfuric acid on cave-wall surfaces causes aggressive carbonate rock dissolution and replacement by gypsum during sulfuric acid speleogenesis. Reddish-brown crusts cover the gypsum in Lower Kane Cave (Wyoming). The crusts are composed principally of C, O, and Si, with abundant microorganisms and euhedral quartz microcrystals. Condensation droplets hang from cave-wall surfaces and average droplet pH was 1.7, ranging from pH 1.25 on crust to 2.92 on gypsum. Droplets on crust had contact angles >90° (avg. 121.6°), indicating hydrophobicity. Droplets with pH < 2 were undersaturated with respect to gypsum, while droplets with pH > 2 were in equilibrium. Condensate solutions on gypsum will approach pH 2, but typically not exceed it, due to buffering by the bisulfate-sulfate weak acid/base pair (pK = 1.92) combined with the gypsum-sulfate. Droplets on crusts had pH values below the critical HSO₄⁻:SO₄²⁻ pK as a result of crust hydrophobicity and acid-producing bacteria. Therefore, armoring of the cave walls by gypsum and biofilms fundamentally impacts how a cave enlarges during sulfuric acid speleogenesis. Microbial colonization of the low pH, moist gypsum habitat forms an organic film that eventually becomes impermeable. Condensation becomes separated and out of equilibrium with respect to the underlying gypsum, precluding diffusion of sulfuric acid through the gypsum to the underlying limestone, limiting or shutting-off sulfuric acid dissolution completely. Only when fresh limestone is exposed will subaerial speleogenesis be reinitiated, and the replacement-colonization cycle start again.

PALEOSOLS & PALEOKARST: A KEY TO PALEOCLIMATE INTERPRETATION IN CARBONATE ISLANDS

V. Ersek, J. Mylroie, J. Mylroie & B. Panuska, Dept. of Geosciences, Mississippi State Univ., Mississippi State, MS 39762; J. Carew, Dept. of Geology, College of Charleston, Charleston, SC 29424

Bahamian paleosols carry a paleoclimatic signal. Paleosol development is dependent on: climate, vegetation, soil organisms, parent material, topography, and time. The Bahamian parent material is Late Quaternary carbonates, and soil developed is therefore linked to epikarst processes. Not only should the paleosol carry a paleoclimatic signal, but the nature of the paleosol/paleokarst contact should also reflect climate: e.g., wetter conditions creating a more complex paleokarst surface that is infilled with more highly weathered paleosol material. Vegetation and soil organisms are a function of climate; the parent material is uniform, so the remaining variables are topography and time. Topography can be factored out by taking sample transects from hills through valleys. Paleomagnetic analysis (secular variation) of the paleosol and geochronology of the parent material can address the time factor. Bahamian carbonates are ~100% CaCO₃, and the insoluble components of the soils are not residual, but are eolian additives as Saharan dust; so the starting materials are known. Bahamian paleosols in some outcrops are developed on a planar, unkarstified carbonate surface, suggesting arid conditions, while others form over a well-developed epikarst surface with significant relief, suggesting wet conditions. Forming soils can fill epikarst voids and harden. Carbonate dissolution removes the weaker parent rock, leaving hardened paleosols, and soil-filled pipes as positive residual structures, inverting topography. Soil infiltra-

tion can infill caves and resist surface destruction; such fills in Cueva del Aleman, Isla de Mona, are at least 1.6 Ma old.

WATER TRACING USING FLUORESCENT DYES TO DELINEATE GROUNDWATER FLOW PATHS IN LEE COUNTY, VIRGINIA

J.E. Fagan, 403 Franklin Drive, Blacksburg, VA 24060

Two groundwater traces using fluorescent dyes were performed in Lee County, VA. The first trace was to determine the hydrologic relationship between 2 caves near the Cedars Natural Area Preserve. Both caves contain populations of the federally listed endangered Lee County Cave Isopod, *Lirceus usdagalun*. The study was part of an investigation by the Virginia Dept. of Conservation and Recreation in support of the U.S. Fish & Wildlife Service recovery plan for *Lirceus usdagalun*. Background fluorescence levels were established at various sampling sites; charcoal samplers were replaced in the stream in Thompson Cedar Cave and at sampling points in nearby springs. Eight ounces of eosine dye were injected into the stream in an unnamed cave 500 m from Thompson Cedar Cave. Ozark Underground Laboratory analyzed the charcoal samplers. Charcoal samplers from Thompson Cedar Cave contained eosine dye, thus demonstrating a hydrologic connection between the 2 caves.

The second dye trace sought to determine the relationship of an actively subsiding sinkhole with nearby springs and wells. Analysis of background charcoal samplers by the investigator using a luminescence spectrometer found no detectable levels of rhodamine WT in sampled springs. During a significant rain event, one pound of rhodamine WT tracer dye was injected into the intermittent stream flowing into the sinkhole. Post-injection analysis of charcoal samplers found dye on one charcoal sampler, thus demonstrating a hydrologic connection between the sinkhole and Jones-Flannery Spring, which is located 200 m south of the sink-point. No connection with other sampled springs or wells was demonstrated.

SEDIMENTATION & POROSITY ENHANCEMENT IN A BREACHED FLANK MARGIN CAVE
L.J. Florea, Dept. of Geology, Univ. of South Florida, Tampa, FL 33559; J.E. Mylroie, Dept. of Geosciences, Mississippi State Univ., Mississippi State, MS 39762; A. Price, Dept. of Geosciences, Univ. of Arkansas, Fayetteville, AR 72701

San Salvador Island, Bahamas, provides a unique location to study modern sedimentation processes on carbonate platforms. The time span of visible geology is highly compressed to the middle Pleistocene through Holocene (< 500 ka). Altar Cave, formed within an oxygen isotope substage 5e eolianite (~125 ka) of the Grotto Beach Formation, is a classic example of a flank margin cave exposed during hillslope retreat. The nature of Altar Cave (restricted entrance, simplistic morphology, and easy access) allowed association of cave features with current environmental conditions and made a sedimentation study easier to perform. Sediment profiles from trenches at 3 locations show that deposits formed in conjunction with a Holocene strand plain present today between the cave and the present beach. ¹⁴C dates show these deposits to be Holocene. Dates from sediment and bedrock from the back of the cave, and XRD and geochemical analyses show the surficial sediment to be recent and that leaching has altered the cave floor bedrock. Petrology of the floor rock has provided the first evidence of autogenic sedimentation in the form of dissolution residuum, most likely drifting downward from the roof, accumulating during void development. Petrographic analyses show that this leaching has resulted in increased bedrock porosity below the soil profile, and introduced organics have contaminated the late Pleistocene bedrock with young carbon, producing a ¹⁴C age of 28 ka. These results demonstrate a potential method of porosity enhancement in young carbonates. These porosity-enhanced zones may have implications on understanding recharge to the fresh-water lens on carbonate islands.

THE USE OF ELECTRICAL GEOPHYSICAL METHODS TO DEFINE GROUNDWATER FLOW PATH IMPACTS IN A KARST AQUIFER

P. Grgich, Univ. of Pittsburgh, Dept. of Geology & Planetary Science, Pittsburgh, PA 15260; R. Hammack, Natl. Energy Tech. Lab., Bruceton Research Center, 626 Cochran Mill Road, Pittsburgh, PA 15236

Pollution and alteration of surface and subsurface hydrology increasingly threaten karst areas. In order to protect these fragile and dynamic watersheds, efficient diagnostic methods must be adapted for use in complex karst settings. The use of surface geophysical techniques to delineate possible flow paths of

clean water in karst aquifers is a new application of existing technology. This study examines stream loss in a small tributary of the Youghiogheny River known as Hoyes Run in Garrett County, MD. The stream bounds the pit of the Deep Creek limestone quarry, operated by Keystone Lime Company. During low flow, the stream abruptly terminates in a swallet, leaving ~100 m of dry streambed. Multiple resistivity profiles using the SuperSting™ R8 System were generated along the zone of loss and compared with results of ground penetrating radar and electromagnetic conductivity profiles in the same location. A dye trace using Fluorescein™ confirmed the flow path of water from the stream into the quarry. Cavities detected in the study were evaluated based on resistivity signal as to their contents (air or sediment filled) to determine which ones might be involved in groundwater transport. The study suggests that surface geophysics coupled with hydrologic and geologic analysis can locate possible flow paths for groundwater in a karst aquifer, even in the absence of obvious karst surface expression. Borehole confirmation is slated before remediation measures are executed.

HYDROGEOLOGIC METHODS USED TO PREDICT CAVERN DEVELOPMENT:

RADERS VALLEY, GREENBRIER COUNTY, WEST VIRGINIA

M. Hall, 5226 Slashwood Lane, Spring, TX 77379

Conventional hydrogeologic mapping techniques have been combined with several analytical methodologies to predict cavern development in Raders Valley, Greenbrier County, WV. The area is dominated topographically by Brushy Ridge anticline and Muddy Creek syncline, which bound this open karst valley. Numerous karst features give clues to cave development in the valley. A model for cavern development was created based on observing cavern development in other areas of Greenbrier County. The hypothesis was that 4 hydrogeologic conditions should be met to develop a large cave in the area: 1) The Hillsdale Limestone-McCrary (shale) Formation contact would need to be exposed to surface runoff; 2) A large area of surface runoff should go underground at the contact (sink); 3) The limestone would need to be fractured where the runoff is captured; 4) Captured surface water would need to flow unobstructed along a fracture or fault to form the cave.

Surface geologic mapping was used to locate the Hillsdale Limestone-McCrary Formation contact and faults. Topographic maps were used to locate sinkholes and determine inflow locations to the cave system. Map calculus techniques were used to locate areas of fractured limestone and faults. Dye traces were compared with other nearby cave-spring dye traces to indicate whether groundwater flow was obstructed. An integrated interpretation was made to predict the location of the cave system on the west side of Raders Valley between Tolers Sinkhole and the Lower Spring. Field walking resulted in finding Middle Earth Cave near the predicted location.

AMERICA'S NATIONAL CAVE & KARST RESEARCH INSTITUTE 2003:

THE GEARING-UP PHASE

L.D. Hose & Z.C. Bailey, National Cave & Karst Research Inst., 1400 University Dr., Carlsbad, NM 88220; L. Land, New Mexico Bureau of Geology & Mineral Resources, Carlsbad, NM 88220; P.J. Boston, Dept. of Earth & Environmental Sci., New Mexico Inst. of Mines & Technology, Socorro, NM 87801

America's National Cave and Karst Research Institute, established by the U.S. Congress, began its "gearing up phase" in December 2002 following a 2.5 year development stage led by Interim Director, Zelda Chapman Bailey. Permanent director Louise Hose has established the Institute office in Carlsbad, New Mexico. The Director's current efforts include initial operational setup, recruiting staff positions, and working with partners to design a permanent building and form a long-term vision.

The Institute operates under a mandate to at least match its current federal funding with non-federal sources. We currently operate under a funding match from the state of New Mexico, which supports collaborative efforts by NM Bureau of Geology-Carlsbad Office hydrogeologist Lewis Land and NM Institute of Mines and Technology (NMT) geomicrobiologist Penny Boston. Land's recent investigations of variations in groundwater discharge from gypsum sinkholes at Bottomless Lakes State Park, NM, has resulted in presentations and a paper to be published this fall in an Oklahoma Geological Survey Circular. NMT created a Cave and Karst Studies Program to provide strong intellectual and educational foundations for the Institute and wider speleological community. Several new Cave and Karst graduate students to be admitted in Fall 2003 under Boston's direction will strengthen NMT's new but aggressive cave and karst research mission.

THE NATIONAL CAVE & KARST RESEARCH INSTITUTE:
HOW WILL IT ADDRESS ITS SCIENCE MANDATE?

L.D. Hose, National Cave & Karst Research Inst., Carlsbad, NM 88220 ; P.J. Boston, New Mexico Inst. of Mines & Technology, Socorro, NM 87801

The U.S. Congress directed the National Cave and Karst Research Institute to pursue 6 objectives, including 3 scientific pursuits: 1) further the science of speleology; 2) centralize and standardize speleological information; 3) foster interdisciplinary cooperation in cave and karst research programs. The enabling legislation directed the National Park Service (NPS) to establish and "jointly administer" the Institute with a public or private agency, organization, or institution. This language, along with a requirement for >50% non-federal cost-sharing, suggests an unfamiliar undertaking for the NPS. The legislation provides ample opportunity for the karst science community to address 2 critical questions about the Institute's science mandate: 1. How will it be achieved? 2. How should the Institute structure itself to achieve it? The Institute's science agenda must be determined before settling on an organizational structure. Current suggestions range from a clearinghouse for grants supporting others efforts to developing full-fledge research facilities and staff at multiple locations.

The NPS "jointly administers" lands with other federal agencies and leads the Cooperative Ecosystem Studies Units Network, examples of multiple, public and private partners doing research. However, both models depend almost entirely on federal funding. A single university (e.g., Los Alamos NL) or corporation (e.g., Sandia NL) operates many federally funded national laboratories under contract. Non-profit corporations commonly administer institutes that need to raise non-federal funds (e.g., Woods Hole). The Institute encourages all karst scientists to join in considering the appropriate science-related activities and most effective organizational structure for the Institute.

ON THE DEFINITION OF CAVERN BREAKDOWN: A REVIEW OF THE CHANGING CONCEPTIONS OF BREAKDOWN

R.A. Jameson, Dept. of Chemistry, Physics & Geology, Winthrop Univ., Rock Hill, SC 29733

Published accounts of cavern breakdown often have equivocated between (1) the use of breakdown as a noun signifying collapsed bedrock fragments, and (2) the use of breakdown as a verb signifying the breaking and falling of bedrock from cave ceilings and walls. A focus on breakdown and breakdown features naturally leads to detailed descriptions of breakdown morphology and passage features generated by the production of breakdown. A focus on collapse processes leads to an emphasis on geologic triggering mechanisms and collapse mechanics, especially if the investigator is concerned with cave stability and safety. The 2 approaches are not mutually exclusive. However, an emphasis on the collapse processes to the exclusion of detailed descriptions of actual, recognizable collapsed bedrock fragments has led to an impoverished conception of cavern breakdown, one that fails to appreciate the richness of breakdown occurrences. One has only to peruse the definitions of breakdown offered in recent lexicons of cave and karst terminology, or offered in putative reviews of breakdown morphology, to appreciate that the time has come for a reassessment of our conception of cavern breakdown. The reassessment should result in a definition that includes a greater variety of phenomena and features, and is applicable to a variety of types of caves, including those formed in carbonates, gypsum, halite, quartzite, sandstone, and lava.

GENESIS & CHARACTERISTICS OF TUMBLING ROCK CAVE, A VALLEY WALL CONDUIT IN JACKSON COUNTY, ALABAMA

P.N. Kambesis, Hoffman Environmental Research Inst., Western Kentucky Univ., Bowling Green, KY 42101; I.D. Sasowsky, Dept. of Geology, Univ. of Akron, Akron, OH 44325

Tumbling Rock Cave is a valley-wall conduit developed along the Cumberland Plateau Escarpment of northeast AL. The cave functions as a drain for Round Cove, a closed depression at the head of Mud Creek Valley. The creek sinks in the floor of the valley, reappears in the cave where it sinks again, and then resurges at a spring near the entrance. The conduit is sub-parallel to, and within, the eastern valley wall, suggesting an origin through stress-relief fracturing. The system is sinuous in plan view, but the orientation of specific segments is controlled by joints, or joint swarms.

Field investigations included cave hydrology and morphology, geochemistry of deposits, and paleomagnetism of clastic sediments to understand the genesis and function of the conduit. Clastic fluvial sediments are found

throughout the cave. These range in size from clay to gravel. Eight oriented samples were collected for paleomagnetic analysis. Six showed normal polarity, suggesting deposition within the present chron. Two samples, taken from a brown clay deposit ~10 m above present stream level, were magnetically reversed. This indicates a minimum age of 780 ka for the growth of this conduit. Conduit growth initiated phreatically and became vadose some time prior to 780 ka. As the cave developed, new levels were formed, and shafts developed concurrently. Escarpment retreat processes presently dominate over valley lowering, possibly due to stratigraphic control of the underlying St. Louis Limestone.

THE CARBONATE ISLAND KARST MODEL AS APPLIED TO ROTA (LUTA), TINIAN, & AGUIJAN ISLANDS, COMMONWEALTH OF THE NORTHERN MARIANA ISLANDS
T.M. Keel, K.W. Stafford, J.E. Mylroie, J.R. Mylroie, Dept. of Geosciences, Mississippi State Univ., Mississippi State, MS 39762; J.W. Jenson, Water & Environmental Research Inst. of the Western Pacific, Univ. of Guam, UOG Station, Mangilao, 96923 GUAM

The main components of the Carbonate Island Karst Model (CIKM) are: 1) enhanced dissolution as a result of fresh water/salt water mixing; 2) glacio-eustatic movement of the fresh-water lens; 3) the influence of local tectonics; 4) eogenetic karst, i.e. karst developed in carbonate rock that has not been buried below the range of meteoric diagenesis; and 5) the model has 4 classifications of carbonate islands: Simple Carbonate (no non-carbonate rocks above sea level), Carbonate-Cover (non-carbonate rocks above sea level but not exposed), Composite (carbonate and non-carbonate rocks exposed), and Complex (inter-fingering of non-carbonate rocks and/or faulting).

Recently, Tinian, Aguijan, and Rota (Luta) in the Mariana Arc were investigated relative to the CIKM. Based on the fieldwork, Tinian is classified as a Composite Island within the CIKM. However, by applying different CIKM classifications to different physiographic regions of Tinian, the Northern Lowland and Median Valley are Simple, the Southeastern Ridge and Central Plateau are Carbonate-Cover, and the North-Central Highland is Composite. Aguijan, a small island just southwest of Tinian, is classified as Simple because it has no non-carbonate rocks exposed, and no evidence of non-carbonate rocks interacting with the lens inside the island. Rota, with large amounts of non-carbonate rock exposed at the Sabana and Talakaya areas, is classified as a Composite Island within the CIKM; however, the Taipingot Peninsula, which is connected to Rota only by a narrow isthmus and has no exposure of non-carbonate rock, could be classified separately as a Simple Carbonate island.

APPLYING STATISTICAL ANALYSIS TO CAVE MAPS TO INFER CONTROLS ON CONDUIT DEVELOPMENT IN THE EDWARDS AQUIFER OF CENTRAL & SOUTH TEXAS

A.L. Lindley, Univ. of Texas at Austin, Dept. of Geological Sciences, Austin, TX 78712; S.D. Hovorka, Univ. of Texas, Bureau of Economic Geology, Austin, TX 78713-8924

As one aspect of a multifaceted approach to a better understanding of flow in karst aquifers, hydrogeologic controls of cave morphology in the Edwards aquifer of central and south Texas were studied. Extensional faulting and fracturing of the Balcones Fault Zone influenced extensive karst development in the Cretaceous Edwards Group limestones. Cave maps from the Texas Speleological Survey archives were digitized using ScionImage Beta Release 4 to collect statistical information on the overall trends of caves. Cave trends were compared to structure defined by mapped faults and gradient to determine the significant controls on flow as reflected in cave elongation. Modern local hydrologic gradient was approximated by assuming it is parallel to topography and measured from DEMs from the Texas Natural Resource Information System.

Structural controls proved to be the main contributor of the morphologies of caves in this region with 23% of 129 caves studied showing strong elongation parallel to the mapped faults. Another 20% of caves were strongly elongated but not parallel to faults or gradient; these were inferred to be developed along local fracture trends. Caves that showed elongation parallel to both structure and gradient comprised 8% of the caves studied. Only 5% of the caves studied were determined to be parallel only to the modern local hydrologic gradient as defined by topography. The remaining 44% of caves could not contribute information toward understanding the phreatic flow system because they are either dominantly vertical caves formed by vadose processes or are weakly elongated phreatic caves.

RECHARGE AREA DELINEATIONS IN THE SALEM PLATEAU, ILLINOIS

P.L. Moss, Ozark Underground Laboratory, 1572 Aley Lane, Protem, MO

Detailed groundwater tracing in the Salem Plateau of southwestern Illinois was conducted from 1996 to 2001. Over 100 dye introductions were made during these investigations. The karst area formed on the St. Louis Limestone in the Salem Plateau physiographic province of Illinois covers ~427 km²; ~120 km² are within recharge areas delineated in these investigations. The study area averages ~29 sinkholes/km² and is autogenic with nearly 100% internal drainage. Studies of groundwater flow in the region have been promoted by those interested in protecting groundwater resources on which rural residents are dependent, and to provide data to support efforts protective of the federally listed Illinois Cave Amphipod (*Gammarus acherondytes*) and its associated biological community. The investigations revealed misunderstandings about the size and interconnectedness of groundwater systems in the region. It also provided an opportunity for coordination with cave bioinventories and a better understanding of cave fauna distribution. Locating potential dye introduction points led to the discovery of many additional caves, several containing species of concern. Six groundwater systems providing habitat for *G. acherondytes* were delineated along with 5 adjacent groundwater systems that recharge delineations were incidental to the investigations. The investigations led to the confirmation that some new locations of *G. acherondytes* represent new populations. Some recharge area delineation has been developed for 4 other groundwater systems. The time of first arrival of dyes and mean groundwater gradients were calculated for each trace.

SEASONAL CHANGES IN SPRING WATER QUALITY IN RUTHERFORD COUNTY, TENNESSEE

A.E. Ogden, C. Kenned & J. Bales, Dept. of Geosciences; R. James, Dept. of Chemistry; J.P. DiVincenzo, M. Rebecca & M. Niese, Dept. of Environmental Science & Technology, Middle Tennessee State Univ., Murfreesboro, TN 37130

Ten successful ground water traces were conducted to 7 springs in Rutherford County, TN. Four of the springs were sampled at week intervals from August 2002 to May 2003. Sampling of 3 additional springs then began in February 2003. The pH of all springs remained fairly constant throughout the sampling period. All springs showed low ammonia and BOD₅ levels. Although the COD levels were also low, they were greater than BOD₅ levels in all instances. This suggests either the presence of nonbiodegradable organics or reduced metals. The results also show that COD levels in the urban-drained springs are higher than rural-drained springs. COD levels could also be correlated with rain events. The highest levels typically occurred during periods of no rain. Dissolved oxygen levels for all springs were lowest during September dry conditions (≤ 5.0 mg/L) and highest after spring rain events. This is expected due to cooling temperatures. DO levels in the urban springs remained below 6 mg/L throughout the sampling period with the lowest level being 2.1 mg/L. Nitrate, phosphate, and sulfate levels were well below health limits but, in general, showed a gradual rise in late spring and early summer. All of the springs showed significant changes in conductivity with storm events. Also, temperature changes of about 5°C were observed between the cold and hot season. This demonstrates open conduit conditions between the sinking streams and springs. As a result, contaminants that enter the karst drainage systems receive little or no filtration before emerging at the springs.

ANNUAL BANDING IN SMALL COLUMNAR STALAGMITES

V.J. Polyak, J.B.T. Rasmussen, & Yemane Asmerom, Earth & Planetary Sciences, Univ. of New Mexico, Albuquerque, NM 87131

Most stalagmites show banding. The frequency of banding in stalagmites is of much interest because they may preserve data on seasonal changes that take place in caves. We are finding well-preserved banding in small columnar stalagmites as couplets of clear and dark (inclusion-rich) layers. These bands are easily observed in typical thin sections. The average thickness of these bands (couplets) matches the uranium series-determined growth rates for the stalagmites, showing that the bands are annual depositions of calcite. Where the banding is continual and not interrupted by growth hiatuses, the band record provides an annual resolution of growth history. The growth history, whether it is represented as growth hiatuses, mineral changes, or an annually resolved history of growth, is likely tied to climate change. The banding record has potential to yield a calibrated annually resolved climate record.

THE EGEMEIER MODEL MEETS HOT AIR: A VADOSE CONVECTIVE AIR-CIRCULATION MODEL FOR THE DEVELOPMENT OF BONEYARD, CEILING PENDANTS, LOFTS, BLIND POCKETS, VENTS, RIMS & SCALLOPS

J.M. Queen, Dept. of Geology, Pomona College, Claremont, CA 91711

Boneyard and associated passage morphologies have traditionally been interpreted as phreatic features. Egemeier suggested that these latter features might result in places from wallrock dissolution by H₂S, dissolved in vadose films, oxidizing to H₂SO₄. But this did not explain why these morphologies developed. A model is proposed whereby convectively driven circulation may develop these morphologies even without H₂S. Atmospheric convection within caves is commonly driven by variations in air density related to depth, temperature, and humidity. Warm, humid air rises and cools. Where it reaches the dew point, continued upward flow results in condensation on exposed surfaces, etching them. This Rayleigh-Benard convection may take place in pores larger than a few decimeters. Irregular recesses in the pre-drainage dissolutional surfaces may be sites of greater heat-flow and consequent cooling, and condensation. They may be selectively deepened, leaving protruding areas to experience progressively less condensation and dissolution. This may result in the development of blind pockets, lofts, domes, pendants, arches, and in places, *en echelon* flow scallops. Where the sinking limb of one convective cell is near the rising limb of another, evaporation gradients result, in which initially acidic condensates are drawn through the intervening bedrock, dissolving some. This may eventually connect the two cells, allowing subsequent barometric pumping of the cave atmosphere. An integrated, connected system of boneyard may result, preserving a sense of the geothermal gradient as a series of vents, dissolved on one side and with a rim of precipitates (popcorn) developed in places on the other.

GUADALUPIAN SPELEOGENESIS REINTERPRETED: NEW MODELS FOR OLD HOLES

J.M. Queen, Dept. of Geology, Pomona College, Claremont, CA 91711

A speleogenetic model, which assumes gypsum blocks, crusts and associated minerals, and dominant passage morphologies are late pre-drainage features, and relies on post-Basin and Range oxidation of H₂S rising from distal hydrocarbon sources is currently popular. However: most gypsum and associated minerals are probably vadose; passage morphologies commonly reflect significant vadose modification; H₂S transport as small bubbles is unlikely in undersaturated water; no evidence of rapid transport is recognized; alternative sources of isotopically light sulfur exist; regional structure favors other gas migration pathways; available dates for cave-lining calcite spar predate the Basin and Range.

An alternative model is proposed: The principal generation of caves initially formed by brine/fresh water mixing following Laramide uplift, unaffected by sulfide oxidation. Continued development resulted in reduced brines filling caves below the water table. H₂S degassed, resulting in reducing atmospheres in caves with limited entrances, and circulated repeatedly by Rayleigh-Benard convection, corroding wall-rock in parts of the rising limbs of cells and precipitating "popcorn" in parts of the descending limbs. Corrosion produced boneyard, domes, arches, rims, etc. Later, as stratigraphic cover thinned and entrances formed, H₂S oxidized to H₂SO₄, etched the wallrock and allowed sulfate minerals to form evaporitically. Condensates separated from bedrock by a gypsum rind may become quite acidic, allowing the local precipitation of acidophilic minerals, including alunite and halloysite, which dates thus indicate a minimum age of draining. Basin and Range uplift and climatic drying opened and dried old caves, while reactivated mixing and degassing continue speleogenesis at lower depths.

CAVING ON THE SAN ANDREAS FAULT: TALUS CAVES IN PINNACLES NATIONAL MONUMENT, CALIFORNIA

B.W. Rogers, US Geological Survey, 345 Middlefield Rd., Menlo Park, CA 94025; J. Despain, Sequoia National Park, CA 93271; W. Frantz, PO Box 2282, Menlo Park, CA 94025 & J. Portillo, Pinnacles National Monument, Paicines, CA 95043

Moderately long talus caves have formed in Pinnacles National Monument, located 180 km south of San Francisco, CA. The Pinnacles are located in the western half of a 23-Ma Miocene volcano torn in half by the San Andreas Fault. The fault has rifted the halved volcano northward nearly 325 km from the Lancaster area, northeast of Los Angeles, to near Hollister in central California. The rhyolitic volcanic rocks lie on a basement of Permian(?) metamorphosed clastic and carbonate sedimentary rocks intruded by Upper

Cretaceous granitic rocks. Three relatively large (1754, 229, and 176 m long) talus caves have developed by both stream erosion along vertical joints sub-parallel to the adjacent San Andreas Fault (SAF) and cyclopean boulders topped by large magnitude ($M > 7$) earthquakes along the SAF. Cave passages range from lofty stream canyons to between-boulder squeezes. Seasonal streams and periodic El Niño floods rearrange cave sedimentary fills, opening and blocking cave passages. Decorations of cristobalite, gypsum, birnessite, ice, and amberat are present. Townsend's Big-eared bats roost and hibernate in different parts of one cave and have staged a minor population boom following closure of the cave following El Niño damage in 1997. Scores of adjacent smaller caves and shelters have formed by tafoni development.

A PRELIMINARY MINERALOGY OF TALUS CAVES IN PINNACLES NATIONAL MONUMENT, PAICINES, CALIFORNIA

B.W. Rogers, US Geological Survey, MS-999, Menlo Park, CA 94025

Moderately long talus caves have formed in Pinnacles National Monument, located 180 km south of San Francisco, CA. Because of the unusual setting, the caves essentially lack "normal" carbonate speleothems. Limited amounts of clear to milky-white colored cristobalite coralloids; biogenic cristobalite diatom moonmilk; gypsum as powdery white crusts and yellow moonmilk, altered by biogenic processes; seasonal ice speleothems; thin, blue-black biogenic deposits of birnessite; and dull to lustrous brown-colored deposits of amberat are present as decorations. The gypsum and cristobalite have formed by evaporation of solutions rich in dissolved silica derived from unstable volcanic ash. The ice forms during the cold winter months from groundwater seepage. The birnessite and cristobalite moonmilk have formed by biogeochemical processes in damp passages. The amberat has formed from woodrat (*Neotoma* sp.) droppings.

QUANTITATIVE FOOTPRINT OF FLANK MARGIN CAVES

M. Roth, K. Zellner & J. Mylroie, Dept. of Geosciences, Mississippi State Univ., Mississippi State, MS 39762

Flank margin caves are hypogenic features resulting from mixing dissolution between freshwater and sea water, and develop with no human-sized openings to the surface. Their morphology is globular chambers that are horizontally broad, and vertically restricted, elongated parallel to the edge of the freshwater lens in which they form. At the back of the main chambers, there are often smaller chambers and small tubes that end abruptly in solid bedrock. This configuration is very different from typical epigenic caves, which are coupled to surface water flow patterns.

This preliminary study examined the "footprint," or quantitative configuration, of flank margin caves. Existing cave maps were analyzed in Autocad to determine cave perimeter and area. Ratios were calculated to determine the correlation between the area and perimeter of the voids in various island settings, taking into account interior residual bedrock pillars. The data reveal that as flank margin caves become larger, they link together laterally along the edge of the lens such that their morphology shifts from circular and sub-oval voids to elongated ovals, a pattern perpendicular to their water flow path. In contrast, epigenic caves grow with an elongation directed parallel to their water flow path.

AQUEOUS GEOCHEMICAL STUDY OF TUFA CREEK, SHANNON COUNTY, MISSOURI

J.A. Schaper & C.M. Wicks, Dept. of Geological Sciences, Univ. of Missouri, Columbia, MO 65211

Carbon dioxide off-gassing due to agitation as Tufa Creek gradually falls 25 m in elevation over a 583-m stream reach is the principal mechanism for calcite deposition along this fen-fed stream. Water chemistry results support this conclusion. The stream water warmed from 14-19°C. The pH increased from 7.02 to 7.96 standard units. Specific conductivity decreased from 0.570 to 0.488 millimhos/cm. Alkalinity decreased from 368-321 mg/L. Dissolved calcium decreased from 157-117 mg/L. Magnesium decreased only marginally from 156-148 mg/L. Total hardness (calcium and magnesium) decreased from 313 to 265 mg/L. Chloride remained statistically unchanged from 7.6 to 7.5 ppm. Sulfate decreased from 5.7 to 4.7 ppm. Calcite saturation increased from -0.03 to 0.85. PCO_2 equilibrated from -1.51 to -2.56, but did not achieve atmospheric (-3.5) before entering the river. Biological mediation perturbs water quality data at a sedge-muck section of the forested fen. Tufa formation correlates with elevation drop. Comparing Tufa Creek data to nearby Ebb and Flow Spring and Thompson Creek reveals that sufficient stream mineraliza-

tion and optimal stream geometry are needed for freshwater calcite deposition. Tufa deposition rates along Tufa Creek are less than those predicted by the Plummer-Wigley-Parkhurst rate law. We speculate this is due to inhibiting effects of dissolved Mg^{2+} or rate changes effected due to periodic gravel scour on the stream.

SUBTERRANEAN SOIL DEVELOPMENT

M.N. Spilde, Inst. of Meteoritics, Univ. of New Mexico, Albuquerque, NM ; P.J. Boston, Dept. of Earth & Environmental Science, New Mexico Tech, Socorro, NM ; D.E. Northup, Dept. of Biology, Univ. of New Mexico, Albuquerque, NM

Low density, soil-like material lines the walls, floors and ceilings of some caves. In Lechuguilla and Spider Caves in New Mexico, this material is particularly well developed, but similar material has recently been described in Jewel Cave, SD. The deposits are diverse in composition with variable amounts of clay and Al-oxide minerals and all are rich in Mn- and/or Fe-oxides. The mineralogy and microbiology of these deposits bear striking resemblance to terrestrial soils in which iron, aluminum and manganese oxides accumulate.

Lithiophorite, nordstrandite (and gibbsite), goethite, kaolinite, and illite have been identified by XRD, and abundant nanocrystalline Fe-oxides were found by TEM examination, similar to minerals found in laterite soils. Likewise, todorokite and birnessite, often observed in soil nodules, are also present in these cave deposits. A diverse microbial community has been identified in these deposits by 16S rDNA sequence analysis, and includes microorganisms whose closest relatives are manganese- and iron-oxidizing bacteria and nitrogen-fixing bacteria.

The similarities in mineral and microbial species composition to that in soils suggest that these cave deposits may undergo a process similar to terrestrial soil development. Both chemical and microbial processes probably influence the formation of these cave deposits. Chemical weathering from the condensation of weak carbonic acid in the cave atmosphere may contribute to the dissolution of the bedrock carbonate, but microbial breakdown of bedrock probably plays a more significant role, as the microorganisms themselves may release organic acids. The residual weathered products, rich in secondary minerals and organic matter, are essentially subterranean soils.

EOGENETIC KARST DEVELOPMENT IN THE MARIANA ISLANDS: AGUIJAN, ROTA, & TINIAN

K.W. Stafford, T.M. Keel, J.E. Mylroie, J.R. Mylroie, Dept. of Geosciences, Mississippi State Univ., Mississippi State, MS 39762; J.W. Jenson, Water & Environmental Research Inst. of the Western Pacific, Univ. of Guam, Mangilao, 96923 GUAM

The islands of the Mariana Arc are tectonically active and composed of Eocene volcanic rocks mantled by Tertiary and Pleistocene carbonates that form an eogenetic karst landscape (i.e., formed on limestones that have not undergone deep-burial diagenesis). Fieldwork on Aguijan, Rota, and Tinian has revealed distinct cave patterns controlled primarily by the position of the fresh-water lens as predicted by the Carbonate Island Karst Model (CIKM), including flank margin (FM) caves and banana holes, and cave morphologies modified primarily by joints, faults, and fractures. Extensive horizons of FM cave development along coastal cliffs and inland scarps, representing previous sea-level stillstands, are present on all 3 islands (e.g., Middle Terrace, Aguijan; As Matmos, Rota; Suicide Cliffs, Tinian), while individual FM chambers can exceed 35 m x 70 m in area and 15 m in height (e.g., Liyang Dangkolo, Tinian). Contacts between non-carbonate and carbonate rock produce closed depressions and show evidence of point source, allogenic recharge (e.g., Lasu Recharge Cave, Tinian; Sabana Recharge Cave, Rota). Fresh-water discharge occurs along carbonate/non-carbonate contacts (e.g., Mattan Hanum, Rota) and along coastlines in the form of seeps and springs (e.g., Gecko Cave, Tinian). Fissure development, primarily as bank-margin failure fractures, has produced caves several hundred meters long and over 30 m deep (e.g., Masalok Fracture Cave Complex, Tinian) by enhanced dissolution along preferential flow paths provided by the fractures, which act as vadose fast flow routes. Tectonic uplift, glacio-eustasy, and cliff margin retreat overprint the various cave features.

A DECADE OF PROCESS-RESPONSE OBSERVATIONS OF A SINKHOLE COLLAPSE CYCLE IN PEBBLE PILE CREEK, LILBURN CAVE, CALIFORNIA

J.C. Tinsley, 1040 Oakland Avenue, Menlo Park, CA 94025; J.W. Hess, 355 Indian Peaks Trail West, Lafayette, CO 80026; D. Cowan, Friars Hill Road, Frankford, WV 24938; H. Hurtt, 623 East Yale, Fresno, CA 93704; W. Farr, 917 Wildrose Avenue, Monrovia, CA 91016

In early 1989, a sinkhole collapsed in the thalweg of Pebble Pile Creek, a tributary to Redwood Creek in the Redwood Canyon karst, Kings Canyon National Park. A compass-and-tape survey of the sink indicated the minimum volume of the collapse to be at least 140-160 m³ of coarse cobbly alluvium. Three sizeable trees up to 0.9 m in diameter were also ingested. A berm of sandy sediment up to 0.3 m thick and largely lacking in fines was found ringing the stream-side of the sink, indicating the collapse was forceful enough to eject streambed sediment. However, the mechanism of ejection is speculative. About 120 m subsurface in Lilburn Cave, the detritus was transported through several conduits by flowing water and reached the South Seas (southern) sump, briefly compromising discharge at the Big Spring stage recorder. Water level at the Z-Room was raised 3-4 m, stabilizing at ~3 m for several years. Progressive in-filling of the sinkhole occurred as the north wall of the sink declined, and the creek transported bedload to the sink during runoff, eventually re-filling the sinkhole, restoring the pre-collapse stream gradient. The 1996-1997 El Nino winter cleared the spring system of its enlarged sediment plug (40 m of head) and restored in-cave sump levels to pre-collapse configurations. These observations reveal an intimate connection between surface events and hydrologic response at Lilburn Cave, and underscore the importance of high-discharge events in the evolution of the Redwood Canyon karst.

ORIGIN AND MINERALOGY OF A TECTONIC RHYOLITE CAVE IN BIG BEND NATIONAL PARK, TEXAS

G. Veni, George Veni & Associates, 11304 Candle Park, San Antonio, TX 78249

Mount Emory Cave is located on the upper flanks of the highest mountain in Big Bend National Park, TX, 2149 m msl. Tectonically formed in the Oligocene Burro Mesa Riebeckite Rhyolite (BMRR), the cave is a series of rifts along stress-release fractures that parallel the topographic contours of a mountaintop ridge. Meter and decimeter scale undulations in the passages and walls occur along variable-scale columnar joints. The cave has a known vertical range of 57.6 m; its lower limits coincide with the base of the jointing. The cave's passages total 151.5 m in horizontal length and generally narrow with greater penetration into the mountain.

X-ray diffraction and scanning electron microscopy was used to quantify the BMRR as about 45% (by weight) quartz and 55% albite feldspar with small amounts of weathered clay and hematite. Analyses were also performed on 3 types of secondary deposits. A sub-millimeter thick water-deposited white coating on many cave walls and breakdown is kaolinite, a weathering product of the feldspar. Pale brown rimstone-like features, generally about 1 cm in length and <1 mm high were mostly quartz, weathered feldspar, and kaolinite. They were water-deposited but their geometry is not always perpendicular to overall water flow patterns probably due to adhesion by the clays. Dark bluish-gray and pale brown coatings along fractures were predominantly weathered feldspar and clay; the darker color is consistent with organic staining. No unusual minerals were found for the geologic setting.

HISTORY

THE ROMANTIC & THE CAVES: JOHN MUIR & THE UNDERGROUND ENVIRONMENT
J.C. Douglas, Dept. of History, Volunteer State Community College, Gallatin, TN 37066

Although not his primary focus, John Muir knew that in many regions caves were an integral part of the natural and human landscape. From his 1867 first ventures into Kentucky caves, his 1869 and 1876 visits to major California caves, to his 1904 tour of the Jenolan Caves in Australia, he was fascinated by the underground environment and its features. He was well aware of spelean variety, as he noted limestone caves, lava tubes and trenches, pot-holes or pits, and ice caves at the bases of glaciers. He was also a careful observer of how cultures and people utilized caves; as living shelters, and hiding places, for American Indians, as social and commercial spaces, and as places of wonder and beauty for Euro-Americans. Muir's ideas regarding caves mirrored his larger attitudes towards nature. In contrast to the prevailing

notion of his day that all aspects of the natural world were potential useable resources, Muir was essentially a romantic who sought beauty, self-discovery, and the presence of God in caves. Implicit in his view of caves was the belief that they were important spaces to be conserved, much like the big trees and high mountains he so valued.

CHARLES DARWIN'S INTEREST IN CAVES

F. Grady, Dept. of Paleobiology, MRC 121 NHB, Smithsonian Institution, Washington, DC 20560

The well known naturalist Charles Darwin noted only a few small sea caves in South America during his nearly 5 year trip around the world in the early 1830s. His correspondence and published writings reveal a knowledge of and interest in various aspects of speleology. Prior to publication of *The Origin of Species*, Darwin requested information about cave adapted species and in *The Origin of Species*, he devoted 2 pages to this subject. His writings show that he was also interested in paleontological and archeological caves. However, poor health after his long voyage prevented any possible cave-related field work.

70 YEARS UNDER THE EARTH

W.R. Halliday, 6530 Cornwall Court, Nashville, TN 37205

During the summer of 1933, my father led me down a pole ladder into Devils Kitchen, a spacious travertine cave and popular tourist attraction in Yellowstone National Park. Half a lifetime later, I found readily recognizable 19th Century stereo views of this cave including 2 by Jay Haynes, celebrated photographer of the park. Originally called "The Mammoth Cave of Wyoming", this cave has long been administratively closed because of supposedly lethal levels of CO₂. Also closed is nearby and possibly larger McCarneys Cave.

With new knowledge of safe exploration techniques in warm, hypercarbic caves and a need for fuller understanding of the depositional speleogenesis of various types of travertine caves, these and similar caves of Yellowstone should be reopened to appropriate scientific study.

LAVA TUBE SYMPOSIUM

MAZE DEVELOPMENT IN HAWAIIAN BASALT CAVE SYSTEMS

D. Coons, 586 E. 9th Rd., Rutland, IL 61358

The geologic literature presents a general perception of basalt tube systems as being relatively short, simple, and ephemeral features. This concept is based primarily on study of lava flow systems located throughout the western continental states of America. Two decades of study beneath 3 active shield volcanoes on the Island of Hawaii have revealed a very different "landscape". Working from >280 km of survey accumulated under Mauna Loa, Kilauea, and Hualalai, a very different pattern begins to emerge. In fact, a large percentage of these tube systems is composed of parallels, braids, or mazes. In a few examples, these maze systems constitute interconnected vertical layers to form some of the most complicated skeins of survey ever documented by the caving community.

CAVES OF THE 1919 FLOW, KILAUEA CALDERA, HAWAII

W.R. Halliday, IUS Commission on Volcanic Caves, 6530 Cornwall Court, Nashville, TN 37205

Nearly 200 caves have been identified in the 1919 "Postal Rift" lava flow in Kilauea Caldera. Only a few of these are traditional lava tube caves, and these are limited to the upper slope of the flow. Despite the current Glossary of Geology definition of "tumulus" as a solid structure, several subtypes of hollow tumuli are readily identifiable here, including dome, sinuous, and transverse types. Individual drained flow lobes and complexes thereof constitute the commonest type of cave in the flow. Some of the latter are measured in hundreds of meters.

Lava rise caves are of 2 subtypes: Circumferential and linear. A spectrum of combined and intermediate forms of all these subtypes is present; a lengthy longitudinal flow lobe cavern is intermediate between ordinary flow lobe caves and lava tube caves. This suggests that simple subcrustal drainage of lava may be the initial phase of speleogenesis of a common type of lava tube cave.

VULCANOSPELEOLOGY IN ICELAND

W.R. Halliday, *IUS Commission on Volcanic Caves, 6530 Cornwall Court, Nashville, TN 37205*

The 10th International Symposium on Vulcanospeleology convened in Iceland in September 2002 with an attendance of about 30 (only 2 Americans). It culminated more than a decade's systematic work by the Icelandic Speleological Society. Iceland is one of the world's great areas for lava tube caves and related volcanic phenomena, and wide varieties were visited during field excursions. Papers all were in English, and covered features in Iceland, the Azores, Australia, Hawaii, Samoa, Japan, Italy, Saudi Arabia, Korea, and elsewhere. Several included stream caves. Other topics were the Great Crack, Kilauea Volcano, HI, a hollow dike in Australia, tree mold complexes in Japan and open vertical volcanic conduits (the world's deepest is just behind the capital city of Reykjavik). Ground penetrating radar has shown a very large extension of the long-celebrated Surtshellir System with no entrance-yet. Field excursions covered caves with especially notable features in both SW and NE Iceland; one crossed the desolate glacier outwash plains of the center of the island where 4x4s are essential. Among the caves visited were Leitharendi, Arnahellir, Stefanshellir, Surtshellir, Vidgilmir, Flughellir, and 2 notable hollow linear tumulus caves containing gently flowing crystal-clear hot water. One of these streaming springs currently is too hot for swimming; the other is just plain wonderful. In contrast were massive ice cascades and other speleothems still present in several caves at the end of summer. New caves still are being found, and much work remains to be done.

LOOKING FOR LAVA TUBES IN COLORADO

D. Medville & H. Medville, *11762 Indian Ridge Rd, Reston, VA 20191*

Unlike other adjacent states to the south and west, the state of Colorado has no known lava tubes. Recent tube-fed basaltic lava flows are rare in Colorado and tubes have not been documented in older flows, although some references to the possibility of tubes exist in the geologic literature. These references described entrances in a welded Oligocene ash flow tuff, and in Miocene basalts on Red Mountain and Flatop Mountain in central Colorado. In the San Luis Valley in south-central Colorado, candidate flows originate on the Los Mogotes Volcano, Volcano de la Culebra, and the State Line shield volcano. Here, indications of the existence of lava tubes in a Pliocene olivine andesite, a Pliocene theolitic basalt (Servilleta Fm.), and the Oligocene Hinsdale Fm., a basaltic lava, have been observed. A number of entrances were seen.

SPELEOTHEMS & LAVA FEATURES IN THE LAVA TUBE CAVES OF EL MALPAIS NATIONAL MONUMENT

V.J. Polyak, *Earth & Planetary Sciences, Univ. of New Mexico, Albuquerque, NM 87131*; P.P. Provencio, *Sandia National Laboratories, Albuquerque, NM 87185*

There are at least 8 tube-forming basalt flows within El Malpais National Monument. The lava tube caves within the Monument contain impressive lava features and numerous speleothems (secondary mineralization). The lava features include benches, gutters, linings, lava stalactites, stalagmites, columns, shelves, grooves, and cinder springs. Lava features provide information regarding the character of lava that flowed through the tubes. The speleothems are predominantly crusts, coralloids, iron-oxide coatings, and moonmilk; however, ice stalactites, stalagmites, draperies, flowstone, and columns are common during winter and spring. Speleothemic crusts, coatings, and moonmilk appear to be more abundant in the older lava tube caves. Not only do some interesting speleothems and minerals form in lava tube caves, but the speleothems have potential to reveal information regarding the timing of tube origin and general climate history. The microbiology related to some of these speleothems may also yield interesting findings.

THE RAPID LOSS OF ICE AT MERRILL ICE CAVE, LAVA BEDS NATIONAL MONUMENT
J. Sowers, *6746 Glen Mawr Ave., El Cerrito, CA 94530*; B. Devereaux, *7605 Aumsville Hwy. SE, Salem, OR 97301*; P. Frantz & B. Frantz, *16345 Englewood Ave, Los Gatos, CA 95032*

Beneath the solid ice floor of Merrill Ice Cave, a large dome-shaped cavity formed within the 2-m thick ice that underlies the floor. How the cavity started and how long it took to grow is not known. The ice cavity was first seen in the fall of 1997, when its roof breached the top layer of the ice floor and the small hole created gave us a view into the 3-m-diameter chamber below. The cavity appears to have formed naturally, the result of air flow in the breakdown

floor beneath the ice causing loss by sublimation. We assume the cavity had been gradually enlarging for many years unseen. Over the past 6 years we have monitored the continued growth of the cavity and simultaneous loss of ice by taking yearly measurements of cavity dimensions and ice floor levels. We have also documented artifacts, primarily trash from visitors, as they became exposed in the walls of ice. The entire 2 m thickness of ice appears to be historical in age. The ice is steadily disappearing and it is unclear whether the air flow through the breakdown will prevent ice from building up significant thicknesses again in the future.

PALEONTOLOGY & ARCHAEOLOGY

ARCHAEOLOGY OF THE KIPUKA KANOHINA CAVE SYSTEM, HAWAII COUNTY, HAWAII

D. Coons, *586 E. 9th Rd., Rutland, IL 61358*

Kipuka is a Hawaiian word that describes an area of land left untouched by the most recent lava flows in the region. These smaller "islands" of ancient land preserve interesting and complex ecosystems on the surface, and often protect fascinating cave systems in the subsurface. The Kipuka Kanohina Cave System comprises one of these highly unusual environments. This prehistoric geologic unit was formed by a long and complex network of tube systems that has now been documented as the second longest basalt cave system in the world.

Many stone implements are still lying *in situ* on the surface of the kipuka. The ancient Hawaiians also extensively utilized the entrance and deep cave areas to collect both water and mineral resources. The cave environment provides constant temperature and humidity, so that an extensive collection of more perishable items has been left behind for the archeological community to investigate. This wealth of material has yet to be studied extensively, or comprehensively catalogued. Preliminary radiocarbon dating of charcoal and wood fragments has begun, raising some very interesting questions.

PLEISTOCENE VERTEBRATES FROM SKELETON CAVE, OREGON

F. Grady, *Dept. of Paleobiology, Smithsonian Institution, Washington, DC 20560*

A small collection of bones and teeth were made from Skeleton Cave, OR in 1928 and 1929. The collection contains about 10 species of mammals. The extinct North American lion, *Panthera atrox*, is represented by 2 nearly complete metapodials and probably fragments of a scapula and humerus. These bones were misidentified in the collection as those of a short-faced bear *Arctodus*. A small horse is represented by 15 complete or fragmentary teeth from a single individual. These teeth most resemble those of *Equus conversidens*, a species previously known from Florida, the Great Plains and the Southwest. The remaining specimens in the collection are those of species from the historical or living fauna of the area.

VERTEBRATE REMAINS FROM NINE SNAKE RIVER LAVA TUBES, IDAHO

F. Grady, *Dept. of Paleobiology, Smithsonian Institution, Washington, DC 20560*; D.A. Hubbard, Jr., *Virginia Speleological Survey, 40 Woodlake Dr., Charlottesville, VA 22901*

Exposed samples of bones and teeth were collected during bioinventories conducted in 2000 from 8 BLM Snake River Plain lava tubes; teeth from an additional BLM lava tube were supplied in 2002. Species identified include: bat (*Myotis* sp.), western big-eared bat (*Plecotus townsendii*), pygmy rabbit (*Sylvilagus idahoenses*), jack rabbit (*Lepus* sp.), yellow-bellied marmot (*Marmota flaviventris*), Great Basin pocket mouse (*Perognathus parvus*), mouse (*Peromyscus* sp.), desert woodrat (*Neotoma lepida*), vole (*Microtus* sp.), northern pocket gopher (*Thomomys talpoides*), wolf (*Canis lupus*), dog or wolf (*Canis* sp.), red fox (*Vulpes vulpes*), grizzly bear (*Ursus arctos*), wolverine (*Gulo gulo*), badger (*Taxidea taxus*), spotted skunk (*Spilogale putorius*), horse (*Equus* sp.). The horse dentition patterns were not similar to any of the more distinctive patterns that are known to represent fossil species. Although none of these species are extinct, the grizzly bear (*Ursus arctos*), wolf (*Canis lupus*), and the wolverine (*Gulo gulo*) are extirpated from this area of Idaho. A skull of a badger, *Taxidea taxus*, is remarkable in that some of its measurements exceed the maximum noted for this species in a standard reference. These remains apparently are of recent age.

These bone deposits were of at least 4 types. Some of the occurrences represent den sites, others were lost individuals that were not able to exit the cave, a few were transplanted by packrats or other animals, and some were not identifiable to a specific type of deposit.