

SEASONAL CONTEXT OF BRISTLY CAVE CRAYFISH *CAMBARUS SETOSUS* HABITAT USE AND LIFE HISTORY

Joshua B. Mouser¹, David C. Ashley², Douglas L. Zentner¹, and Shannon K. Brewer^{3,4,C}

Abstract

Cave crayfishes are important members of groundwater communities, but many cave crayfishes are threatened or endangered. Unfortunately, we lack basic life history and ecological data that are needed for developing conservation plans for most cave crayfishes, especially the role of seasonal and annual fluctuations in structuring populations. Therefore, we determined the seasonal life history and habitat use of *Cambarus setosus* in Smallin Civil War Cave, Christian County, Missouri, United States. We conducted visual crayfish surveys over a 400 m section of the cave from 2006 to 2019. We used multinomial logit, multiple linear regression, and logistic regression models to estimate crayfish substrate, water depth, and water velocity use, respectively. All models included sex, carapace length, season, distance into the cave, and interactions between all variables and sex as predictor terms. We also used *t*-tests to assess morphometric differences between male and female crayfish. Six mark-recapture events (2010 to 2019) were used to estimate population sizes using a nil-recapture model. We attempted to age eight individuals using gastric mill bands, but annual bands were not discernable. We found reproductively active males during all seasons. We captured one ovigerous female during the spring, though ovigerous females were observed during show cave tours during spring, summer, and autumn. Male *C. setosus* were more likely to use homogenous and heterogeneous rock substrates and shallower and calmer water when compared to females; however, these relationships varied based on distance into the cave and season. Females sampled were significantly larger than males, and males regenerated chelae more often. Minimum population size estimates ranged from 9 to 159 individuals and indicated the population was relatively stable. Our data provide both a baseline population estimate for comparison with future studies and valuable trait information that is often lacking but useful for developing conservation efforts.

INTRODUCTION

There is broad recognition that cave crayfishes play an important role in groundwater ecosystems, and many populations are at risk of extinction. Crayfishes are keystone species that shape the structure and function of aquatic ecosystems (Paine, 1969). For example, crayfishes serve as aquatic nutrient cyclers (Momot, 1995) and are food for many species (e.g., >200 in the Ozark Highlands ecoregion (DiStefano, 2005)). In groundwater systems specifically, cave crayfishes are part of stygobiont communities that support clean water that is used for drinking and crop irrigation (Danielopol and Griebler, 2008; Boulton et al., 2008; Griebler et al., 2014). Cave crayfishes typically have narrow distributions (Larson and Olden, 2010) and K-selected life histories (e.g., long life span (Venarsky et al., 2012)), resulting in an intrinsically high risk of extinction. For example, approximately 70 % of stygobiont crayfishes are at risk of extinction (Taylor et al., 2007). The persistence of cave crayfish populations is threatened by water pollution, recreational caving, and invasive species (Graening et al., 2006; Mouser et al., 2019).

Effective conservation and management strategies for cave crayfish populations requires an understanding of their life history and habitat use (Moore et al., 2013; DiStefano et al., 2016; Taylor et al., 2019). We lack a basic understanding of both the biology and ecology of many crayfishes, especially cave-dwelling species (Taylor et al. 2019). The lack of basic biological and ecological knowledge is a major impediment in developing meaningful conservation efforts for subterranean organisms (Mammola et al., 2019). Life history data (e.g., age, fecundity, and recruitment) can be used to predict at-risk and invasive crayfish species (Larson and Olden, 2010), to help managers determine appropriate sampling techniques (Crandall, 2016), and to develop habitat restoration strategies that target life-stage requirements (Dyer et al., 2016). Management decisions also benefit from an understanding of a species' habitat requirements. For example, species-habitat associations are particularly useful to help direct restoration efforts (Smith et al., 1996), to control invasive crayfishes (Light, 2003), and to determine potential reintroduction sites (Renai et al., 2006).

The bristly cave crayfish *Cambarus setosus* is the most common cave crayfish of the Ozark Highlands ecoregion. *Cambarus setosus* has been documented at 48 sites (i.e., caves, wells, or springs) in Missouri and two sites in Arkan-

¹Oklahoma Cooperative Fish and Wildlife Research Unit, Oklahoma State University, Stillwater, OK 74078

²Department of Biological Sciences, Missouri Western State University, Saint Joseph, MO 64507

³U.S. Geological Survey, Oklahoma Cooperative Fish and Wildlife Research Unit, Stillwater, OK 74078

⁴Current address: U.S. Geological Survey, Alabama Cooperative Fish and Wildlife Research Unit, 203 Swingle Hall, Auburn University, Auburn, AL 36849

^CCorresponding author; skb0064@auburn.edu.

sas with 164 individuals being reported from all sites (Graening et al., 2006). *Cambarus setosus* is currently listed as stable by the American Fisheries Society (Taylor et al., 2007), near threatened by the International Union for Conservation of Nature (DiStefano et al., 2021), and vulnerable by NatureServe (NatureServe, 2009) and by the Missouri Department of Conservation (Missouri Natural Heritage Program, 2021). Gardner (1986) noted that *C. setosus* was more abundant in stream sections characterized by a sandy, muddy substrate with scattered breakdown and abundant bat guano. Similarly, Marquart (1979) found *C. setosus* in silty substrates with rock, gravel, and organic debris. *Cambarus setosus* can reach 120 mm total length (TL); males can be reproductively active (i.e., form I) at 53 mm TL, and there is little difference in size between the sexes (Pflieger, 1996).

The previous studies of *C. setosus* habitat use and life history provide important insight but are mostly qualitative observations or short-term studies limited to a few caves (but see Marquart (1979)). These studies do not capture seasonal or annual changes within a cave and these dynamics (e.g., changing water levels or flow) may be quite important in structuring the life history and ecology of cave organisms (Jegla and Poulson, 1970; DiStefano et al., 2020). Therefore, the goal of our paper was to provide basic population information for *C. setosus* in a seasonal context. We specifically focused on basic life history (i.e., reproductive timing and size), habitat use (i.e., substrate, water depth, and water flow) and estimating minimum population size of *C. setosus*. Collectively, these metrics are useful for determining growth, mortality, and recruitment (Panfili et al., 2002) and can be used to protect or restore key habitat within caves.

METHODS

Study Area

We focused our study on *Cambarus setosus* within Smallin Civil War Cave (Smallin Cave), Christian County, Missouri, US. Smallin Cave is a show cave located within the Ozark Highlands ecoregion. The Ozark Highlands ecoregion is characterized by karst topography (Unklesbay and Vineyard, 1992), moderate climate (e.g., rainfall of 97–122 cm and average temperature of 13–16 °C (Adamski, 2000)), and mixed forest interspersed with lowland agricultural use (Woods et al., 2005). Smallin Cave has a handicap-accessible tour route elevated 1–2 m above the cave floor, which extends approximately 200 m into the cave from the entrance. From the end of the tour route, the cave extends an additional 700 m. A shallow stream with occasional pools meanders along the floor of the cave. Smallin Cave is home to one of the largest known populations of *C. setosus* with at least 47 individuals (Graening et al., 2006) and a relatively small population of bats (D.C. Ashley, personal observation).

Crayfish and Habitat Surveys

We conducted 38 visual surveys from November 19, 2006 to June 6, 2019, and we also report some opportunistic observations that were noted during cave tours in 2010. We sampled during spring (March–May, $n = 18$ surveys), summer (June–August, $n = 8$), autumn (September–November, $n = 5$), and winter (December–February, $n = 7$). Each survey covered approximately 400 m of the cave and consisted of 2 to 6 observers walking slowly from the cave entrance to the end of the study area visually searching for crayfish on the substrate. The remaining cave passage beyond the study area is a tight crawl passage and was not surveyed. We captured crayfish via hand nets, recorded distance into the cave from the dripline (i.e., cave entrance), and placed a 900 cm² frame at the capture location to estimate microhabitat use. Water depth (± 1.0 cm) was measured in the center of each occupied grid and water velocity was visually estimated as: calm ($n = 161$), slow flow ($n = 51$), moderate flow ($n = 17$), or fast flow ($n = 4$). We combined slow, moderate, and fast flows into a single category (flowing) because these data were naturally bimodal when compared to calm water. Substrate was visually estimated as the proportion of the quadrat comprising mud/silt (particle diameter < 0.01 cm), sand (0.01–0.2 cm), pebble (> 0.2 –6 cm), cobble (> 6 –20 cm), large rock (> 20 –26 cm), or bedrock (> 26 cm). For each crayfish, we measured carapace length (± 1.0 mm) and length of both chelae (± 1.0 mm) using a ruler. We also recorded sex and whether crayfish were reproductively active (i.e., form I males or ovigerous females) or not (i.e., form II males or non-ovigerous females) using visual observation. Males were considered form I if the tip of the gonopod was corneous and pointed, and females were considered ovigerous if they were carrying eggs on their telson. The cave manager also recorded ovigerous females during show cave tours in 2010. On six surveys, a permanent marker was used to label the carapace with a unique number and a repeat survey was conducted 2–3 days later to count the number of marked and unmarked individuals (Table 1). We chose to use permanent marker because it is effective and safe for short-term recapture events (Ramalho et al., 2010) while being cheaper than other methods. Distance into the cave, habitat data, and crayfish morphometrics were not collected during most of the six repeat surveys.

We condensed the substrate estimates into four categories that we hypothesized to be ecologically relevant. We classified quadrats comprising 100 % bedrock, sand, silt, or clay as “bedrock/fine” because they serve as poor shelter for a crayfish (i.e., crayfish cannot burrow into bedrock and too much fine substrate results in suffocation; (Dyer et al., 2015)). Quadrats comprising a mixture of fines and other substrates were categorized as heterogeneous fine because excess fine substrates can suffocate crayfish (Dyer et al., 2015); however, this substrate would present usable shelter,

and *C. setosus* has been shown to use this habitat (Marquart, 1979; Gardner, 1986). All substrate mixtures containing size distributions larger than sand, but excluding bedrock, were classified as “homogenous rock.” Lastly, we classified quadrats with a mix of bedrock and homogenous rock as “heterogenous rock.”

Analyses

We chose variables hypothesized to influence habitat use as predictor terms in our models (described in the next three paragraphs). Crayfish habitat use often differs depending on sex and season (e.g., DiStefano et al., 2013) and size (e.g., Dyer et al., 2016). Therefore, we hypothesized *C. setosus* habitat use depended on sex (female or male), size (i.e., carapace length (CL)), and season (i.e., spring, summer, autumn, or winter). Distance into the cave was included as a predictor term because habitat visibly changed throughout the cave. Lastly, we included interaction terms between sex and all other variables to account for possible differences in habitat use between males and females. Our analyses were conducted using the statistical software R (R Core Team, 2020), and $\alpha \leq 0.1$ was chosen *a priori* as our cutoff for significance.

We built a multinomial logit model to estimate the probability that crayfish were found in each of our four substrate categories (i.e., bedrock/fine, heterogeneous rock, homogenous rock, and heterogeneous fine). Sex, CL, season, distance into the cave, and interactions between sex and all other variables were included as predictor terms. Sex and season were treated as categorical variables with female and spring as reference categories. Our response variable was the probability that crayfish were found in each of the four substrate categories. We used the multinom function within the nnet package (Venables and Ripley, 2002) to fit our model as described by Faraway (2005). We started with the full model and removed predictor terms one at a time that most decreased Akaike information criterion adjusted for a small sample size (AICc; Burnham and Anderson, 2001) until removing terms no longer decreased AICc (Faraway 2005). We assessed model fit using a χ^2 test to compare the observed probability that crayfish were found in each substrate category to the probability predicted by the model (Yau, 2013).

We built a multiple linear regression model to predict water depth used by bristly cave crayfish. Sex, CL, season, distance into the cave, and interactions between sex and the other variables were included as predictor terms in the model. Sex and season were treated as categorical variables with female and spring as reference categories. Our response variable was water depth used by the crayfish. Water depth was natural-log transformed due to a right-skewed distribution (Zar, 1999). Two crayfish were found out of the water and those data were removed due to high influence on the model (i.e., Cook’s distance > 0.5). We selected the best model using the same approach described above. Model assumptions and fit were determined via visual analysis of the quantile-quantile plot (Q-Q plot) and plotting residual versus predicted values (Freund and Wilson, 2003).

We built a logistic regression model to determine whether crayfish were more likely to be found in calm or flowing water. Sex, CL, season, distance into the cave, and interactions between sex and the other predictor variables were included in the model. Sex and season were treated as categorical variables with female and spring as reference categories. Our response variable was calm (0) or flowing water (1). We selected the best model using the selection approach previously described. We assessed model fit using binned residual plots because traditional residual plots are uninformative for models with binary response variables (Gelman and Hill, 2007).

We assessed morphometric and gender differences using *t*-tests ($\alpha \leq 0.1$). We used Welch’s *t*-tests to compare groups with unequal variances as indicated by an *F*-test. We used unpaired *t*-tests to determine if there were significant differences in male and female crayfish for both average CL and average left and right chelae size divided by CL (i.e., adjusted for overall size). We hypothesized that a larger difference in left and right chelae size would reflect loss and partial chela regeneration resulting from aggressive behavior. Therefore, we also compared the left and right chelae size via paired *t*-tests for both males and females.

A nil-recapture model was used to estimate the minimum expected population size of bristly cave crayfish in Smallin Cave. The nil-recapture model was selected because recaptures during each event ranged from 0 to 2 and because Smallin Cave can be classified as an environment that is difficult to sample (Bell, 1974; Friedenberget al., 2018). Using Bayesian methods improves the interpretation of the resulting model by allowing statements about the probability that abundance exceeds some lower bound. Because spatial data were not available, the spatial distribution of crayfish was assumed uniform and population estimates were obtained from the posterior gamma distribution where the shape and rate parameters were defined using:

$$\alpha = \alpha_0 + R$$

$$\beta = \beta_0 + \sum_{t=1}^{i-1} c_t m_t$$

where α is the shape parameter defining gamma distribution, α_0 is the prior for the shape parameter, R is the total number of recaptured individuals, β_0 is the rate parameter defining gamma distribution, c_t is the prior for the rate parameter,

is the number of individuals captured in sample event i , m_i is the estimated number of individuals marked during sample event i , and l is the maximum number of sample events (Friedenberg et al., 2018).

This formulation of the nil-recapture model allows an estimate of minimum expected population size (Edwards, 1974) at various probability levels using the reciprocal obtained from the gamma quantile function. To provide a range of minimum expected population size, we *a priori* selected probability levels of 0.99, 0.95, and 0.75. This allowed us to be 99 %, 95 %, and 75 % confident, respectively, that the actual population was greater than or equal to our estimates. Every estimate was obtained using both uniform and Jefferys priors as they are convenient priors that produce equivalent estimates to lower bounds from frequentist and likelihood approaches, respectively. The use of two different priors also allowed us to better understand the effect of prior choice on our minimum expected population size estimates given our limited number of recaptures (Friedenberg et al., 2018).

Crayfish aging

We attempted to age *C. setosus* via the gastric mill to determine the longevity of the species following Mouser et al. (2020). Eight *C. setosus* that died naturally in Smallin Cave during summer 2018 were collected. We extracted the gastric mills, thinly sectioned them, and mounted them on microscope slides. We aged sections from multiple ossicles for each crayfish because the recovered crayfish were in various states of decomposition and not all of the ossicles could be located. The best section from each zygo-cardiac ossicle and pterocardiac ossicle, and the two best sections from the mesocardiac ossicle were mounted on each slide. Two readers attempted to age the slides together.

RESULTS

Cambarus setosus CLs ranged 3.0–45.0 mm ($n = 399$, mean = 22.7 ± 7.2 mm) and were captured during all seasons from a variety of habitats (Table 1). Location of capture ranged between 23–420 m from the dripline. Both male ($n = 167$, mean CL = 22.0 ± 6.0 mm) and female ($n = 133$, mean CL = 24.7 ± 7.5 mm) crayfish were collected during all seasons. Reproductively active males were collected during spring ($n = 8$), summer ($n = 6$), autumn ($n = 5$), and winter ($n = 7$), whereas a single ovigerous female was captured during the spring; all other crayfish collected were not reproductively active. However, ovigerous females were observed during cave tours in May, June, July, and November. The smallest male had an 18 mm CL and the smallest ovigerous female had a 30 mm CL. Crayfish were observed using a variety of microhabitats characterized by different substrate compositions: bedrock/fine substrate ($n = 31$), heterogeneous fine ($n = 31$), heterogeneous rock ($n = 28$), and homogenous rock ($n = 142$). Crayfish were found in 0–106 cm of water (mean = 20.4 ± 17.8 cm) and more often in calm water ($n = 161$) than flowing water ($n = 72$).

The results of the multinomial model indicated that distance into cave and sex were predictors of substrate use (Fig. 1 and Table 2). Male *C. setosus* were more likely to use homogenous and heterogeneous rock substrates than females, whereas female *C. setosus* were more likely to use bedrock/fine substrate than males. The magnitude of the difference was greater near the entrance of the cave for bedrock/fine and homogenous rock substrates. Results of the χ^2 goodness-of-fit test indicated multinomial model fit was appropriate ($\chi^2_6 = 8$, $p = 0.24$).

The results of the multiple linear regression model indicated that water depth use was related to season and distance into the cave, depending on sex (Fig. 2 and Table 3). Male *C. setosus* were more likely to use deeper water than females near the entrance of the cave, but more likely to use shallower water than females farther in the cave. Crayfish were found in shallower water more often in autumn compared to spring. The Q-Q plot and the residual plot showed no concerning trends, suggesting adequate linear model fit.

Similar to the multiple linear regression model, the results of the logistic regression model indicated that water depth use was related to season and distance into the cave depending on sex (Fig. 3 and Table 4). Males were more likely than females to use flowing water near the entrance and less likely to use flowing water farther in the cave compared to females. Crayfish were less likely to use flowing water in the summer and autumn (i.e., negative relationship) compared to spring, when high flows are more common. The binned residual plot indicated good generalized linear model fit because 95 % of the binned residual were contained in theoretical error bounds, and the plot did not reveal any concerning trends.

We also found morphological differences between male and female *C. setosus*. Female crayfish were significantly larger than males ($t_{243,69} = 3.29$, $p < 0.01$). There was not a significant difference between male and female crayfish chelae size when adjusted for carapace length ($t_{286} = 0.31$, $p = 0.76$). Right and left chelae were not significantly different for male ($t_{157} = 1.09$, $p = 0.60$) or female crayfish ($t_{129} = -0.41$, $p = 0.68$). Although more males had regenerated or missing chelae ($n = 37$) than females ($n = 29$), this represented 22 %.

Our minimum population estimates of *C. setosus* were generally low but appear relatively stable through time (Fig. 4). The largest variability in minimum population size estimates was observed when a Jeffreys prior and a probability of 0.75 were used to estimate the minimum population size (range = 32–159). The smallest variability in minimum population size estimates was observed when a uniform prior and a probability of 0.99 was used to estimate the minimum population size (range = 9–23). The Jeffreys and uniform priors gave similar minimum population size estimates when

Table 1. Crayfish counts from visual surveys in Smallin Civil War Cave, Christian County, Missouri, United States. Female crayfish (Fem; $n = 133$) captured ranged from a minimum (Min) carapace length of 7 mm to a maximum (Max) of 40 mm. Male crayfish ($n = 167$) ranged from 8–45 mm. Sex and length were not determined (ND) for some crayfish (Unknown, $n = 91$).

Trip	Date	Female			Male			Unknown			Total
		Count	Min	Max	Count	Min	Max	Count	Min	Max	
1	19 Nov 2006	6	12	33	2	11	20	2	10	12	10
2	28 Jan 2007	4	7	21	5	15	24	2	ND	ND	11
3	18 May 2007	2	18	19	4	19	34	0	6
4	18 Nov 2007	2	19	27	6	14	30	3	3	ND	11
5	29 May 2008	3	27	40	3	12	30	0	6
6	20 Jul 2008	4	23	40	1	26	26	2	ND	ND	7
7	02 Nov 2008	5	21	28	4	28	35	2	ND	ND	11
8	15 Feb 2009	3	24	31	0	0	3
9	28 May 2009	5	23	40	1	33	33	0	6
10 ^a	24 May 2010	5	20	28	3	26	30	4	ND	ND	12 (11)
11 ^b	27 May 2010	3	22	35	2	10	13	3	ND	ND	8 (6,1)
12	28 May 2010	3	31	35	5	19	30	0	ND	ND	8
13	14 Aug 2010	4	7	19	6	12	30	2	ND	ND	12
14	14 Nov 2010	1	26	26	1	19	19	2	ND	ND	4
15	30 May 2011	0	1	19	19	0	1
16	19 Dec 2011	2	24	25	4	21	38	2	ND	ND	8
17 ^a	23 May 2012	7	7	27	14	10	43	5	ND	ND	26 (26)
18 ^b	25 May 2012	2	ND	ND	4	ND	ND	0	6 (6,1)
19	03 Jan 2013	4	15	33	4	15	24	1	15	15	9
20 ^a	22 May 2013	7	25	35	1	35	35	0	8 (8)
21 ^b	24 May 2013	0	4	20	25	1	ND	ND	5 (5,0)
22	21 May 2014	6	10	30	5	16	29	2	ND	ND	13
23	24 Jul 2014	5	12	26	9	14	25	5	ND	ND	19
24	16 Jan 2015	0	5	14	29	2	ND	ND	7
25	18 Jan 2015	2	22	25	2	19	33	3	ND	ND	7
26 ^a	20 May 2015	7	15	30	8	12	30	2	ND	ND	17 (15)
27 ^b	22 May 2015	1	ND	ND	5	ND	ND	4	10 (7,0)
28	25 Ma. 2016	4	25	32	4	16	27	4	ND	ND	12
29 ^a	16 May 2016	4	20	34	6	20	26	0	10 (10)
30 ^b	18 May 2016	1	37	37	0	12	ND	ND	13 (13,1)
31	26 Aug 2016	7	13	40	9	8	26	1	ND	ND	17
32	27 Feb 2017	3	29	32	8	12	30	0	11
33	24 May 2017	1	16	16	1	30	30	0	2
34	02 Aug 2017	8	17	29	5	21	30	1	ND	ND	14
35	26 June 2018	5	19	30	9	16	30	9	ND	ND	23
36	27 Nov 2018	4	11	20	7	18	28	7	ND	ND	18
37 ^a	04 Jun 2019	3	18	45	9	16	29	3	ND	ND	15 (12)
38 ^b	06 Jun 2019	0	0	13	ND	ND	13 (13,2)

¹ Crayfish were marked on this trip. Number in parentheses indicate the number marked.

² Crayfish were recaptured on this trip. Numbers in parentheses indicate the number of crayfish captured (first number) and how many of those were previously marked (if a second number is provided).

using a probability of 0.95 (Jeffreys = 17–55, uniform = 13–35) or 0.99 (Jeffrey's = 12–32, uniform = 9–23). The range of minimum population estimates (i.e., 10–32) during the initial mark-recapture period (May 2010) overlapped with the range of minimum population estimates (19–47) in the final mark-recapture period (June 2019), demonstrating relative stability through time.




Figure 1. Predicted probabilities from our multinomial model of bristly cave crayfish *Cambarus setosus* using different substrate types in Smallin Civil War Cave, Christian County, Missouri, United States. A significant interaction between sex and distance into the cave indicates that male (dashed line) and female (solid line) crayfish have differing habitat uses depending on location within the cave.

Table 2. Mean and standard error (SE) estimates for variables included in the top multinomial model used to predict the probability of substrate category used by bristly cave crayfish *Cambarus setosus* in Smallin Civil War Cave, Christian County, Missouri, United States. Sex and season were treated as categorical variables with female, and spring as reference categories, respectively. Distance (± 1 m) was a continuous variable representing the distance a sample was taken from the cave entrance. Probability of substrate use was modeled with respect to bedrock/fine substrate and parameter estimates of the environmental variables are given for each of the other categories.

Parameter	Mean	SE
Heterogenous fine-intercept	-0.88	0.91
Heterogenous rock-intercept	0.11	0.99
Homogenous rock-intercept	0.86	0.79
Heterogenous fine-male	1.05	0.72
Heterogenous rock-male	1.44	0.73
Homogenous rock-male	1.27	0.62
Heterogenous fine-distance	5.72×10^{-3}	4.92×10^{-3}
Heterogenous rock-distance	-2.03×10^{-3}	5.71×10^{-3}
Homogenous rock-distance	3.38×10^{-3}	4.48×10^{-3}

anism is unknown, the lack of yearly bands may be due to the absence of seasonal changes in temperature that typically influence the formation of growth marks on hard structures (Wright et al., 2002), including gastric mills (Leland et al., 2015; Mouser et al., 2020). Caves have relatively stable temperatures; therefore, the absence of yearly bands may be due to constant growth in those systems or extremely limited growth occurring over much shorter intervals. Limited growth over shorter intervals may also explain the presence of sub-yearly bands, as sub-yearly rings have been observed on fish otoliths due to feeding changes (Wright et al., 2002). In contrast to our findings, Mouser et al. (2019) found that epigeal *F. neglectus* in caves still displayed bands that seemed to reflect annual conditions, but crayfish

We used gastric mill sections in an attempt to age eight *C. setosus*, but our results were inconclusive. The carapace length of the aged crayfish ranged from 13.0–31.0 mm (mean = 19.1 ± 5.7 mm). None of the crayfish displayed clear yearly growth bands (Fig. 5). However, many gastric mills displayed hypothesized sub-yearly bands (Fig. 5).

DISCUSSION

In this study, we provide detailed information on the seasonal habitat use and life history of *C. setosus*. Caves are relatively stable environments in some ways (e.g., temperature); however, they can also be quite dynamic (e.g., flooding) (Simon, 2012). Many studies have shown that seasonal dynamics are important for structuring how cave ecosystems function (e.g., food flux during floods (Poulson, 2012), controlling life history of stygobionts (DiStefano et al., 2020)). Despite this knowledge, most data for *C. setosus* are limited to a short time frame. We found that seasonal dynamics are important for the reproduction and habitat use of *C. setosus* and may affect the formation of gastric mill bands.

Cambarus setosus reproduction appears to follow seasonal patterns, but shifts in seasonal patterns (e.g., rainfall) may alter reproduction timing. We found that crayfish reproduction in Smallin Cave occurred primarily in the spring and early summer (i.e., when the majority of ovigerous females were found); however, form-I males were found during all seasons. These results are similar to reproductive patterns found for surface crayfishes and other cave crayfishes. For example, many species of lotic crayfish in the Ozark Highlands ecoregion molt twice annually for reproductive purposes and lay eggs during the spring (Pfleger, 1996). Less is known about the reproductive habits of cave crayfishes; however, it appears cave crayfish or surface-dwelling crayfish that invade caves may reproduce more often under some circumstances and changes in seasonal patterns may shift reproductive timing. For example, Mouser et al. (2019) found that surface crayfish reproduced all year in a cave; however, most reproduction occurred during spring. Jegla (1966) observed that *Orconectes pellucidus inermis* also followed reproductive patterns similar to surface species; however, flood events were noted by Jegla and Poulson (1970) to shift reproductive timing. Similarly, DiStefano et al. (2020) observed ovigerous *O. stygocaneyi* in August a few months after heavy rainfall. Shifts in seasonal environmental patterns may explain why we found reproductively active crayfish during seasons not typically associated with reproduction.

We did not observe hypothesized yearly bands on *C. setosus* gastric mills, which is contrary to work on *Faxonius neglectus* found in surface streams (Mouser et al., 2020) and caves (Mouser et al., 2019). Although the mech-




Figure 2. Relationship between water depth use by bristly cave crayfish *Cambarus setosus* and distance into Smallin Civil War Cave, Christian County, Missouri, United States. A significant interaction between sex and distance into the cave indicates that male (dashed line) and female (solid line) crayfish have differing habitat uses depending on location within the cave.




Figure 3. Predicted probability of flowing water use by bristly cave crayfish *Cambarus setosus* in Smallin Civil War Cave, Christian County, Missouri, United States. A significant interaction between sex and distance into the cave indicates that male (dashed line) and female (solid line) crayfish have differing habitat uses depending on location within the cave.

Table 3. Mean and standard error (SE) estimates along with *p* values for variables included in the top multiple linear regression model used to determine the association between bristly cave crayfish *Cambarus setosus* and water depth in Smallin Civil War Cave, Christian County, Missouri, United States. Sex and season were treated as categorical variables with female and spring as reference categories. Distance (± 1 m) was a continuous variable representing the distance of the sample from the cave entrance.

Parameter	Mean	SE	<i>p</i> value
Intercept	2.52	0.28	<0.01
Male	0.45	0.34	0.19
Distance	1.71×10^{-3}	1.42×10^{-3}	0.23
Summer	0.10	0.17	0.53
Autumn	-0.42	0.22	0.06
Winter	-0.29	0.21	0.18
Male \times distance	-3.50×10^{-3}	1.88×10^{-3}	0.06

Table 4. Mean and standard error (SE) estimates along with *p* values for variables included in the top logistic regression model used to predict the association between bristly cave crayfish *Cambarus setosus* and flowing water in Small Civil War Cave, Christian County, Missouri, United States. Sex and season were treated as categorical variables with female and spring as reference categories. Distance (± 1 m) was a continuous variable representing the distance of the sample from the cave entrance.

Parameter	Mean	SE	<i>p</i> value
Intercept	-1.29	0.72	0.07
Male	1.41	1.07	0.19
Distance	5.69×10^{-3}	3.68×10^{-3}	0.12
Summer	-0.96	0.49	0.05
Autumn	-2.28	1.08	0.04
Winter	0.14	0.50	0.29
Male \times distance	-0.01	6.79×10^{-3}	0.03

movement out of the cave or food fluctuations derived from maternal colonies of gray bats *Myotis grisescens* may be contributing factors. Using the gastric mill to age cave crayfish could be valuable because it is difficult to determine the longevity of cave crayfish through traditional methods (Venarsky et al., 2012). However, this technique might not be useful for populations with low abundances unless natural mortality is observed. Further, more laboratory work needs to be completed on band formation before the technique could be considered for subterranean crayfishes.

Male and female *C. setosus* are found in different habitats depending on the location within the cave. The observed differences in habitat use between the sexes could be explained by more dominant males excluding females from ideal habitats (Fero and Moore, 2008). However, chelae size was not significantly different in male and female crayfish, but other forms of behavior (e.g., pheromone releases (Schneider et al., 1999)) may cause females to avoid males. Females were typically larger than male crayfish, and body size can influence cover use (Streissl and Hödl, 2002; Dyer et




Figure 4. Minimum expected population estimates from our nil-recapture model for the bristly cave crayfish *Cambarus setosus* in Smallin Civil War Cave, Christian County, Missouri, United States. Points represent the minimum population estimate when there is a 75 %, 95 %, or 99 % chance that the true population is greater than or equal to the population estimate when using a Jefferys (gray shapes) or uniform (black shapes) prior.

al., 2016). For example, male *C. setosus* may avoid faster water because smaller crayfish are more likely to be swept downstream (Hobbs, 1978; Caine, 1978). Female crayfish could be associated with bedrock because they are too large to take refuge under other substrates and crevices in the bedrock that may provide cover.

Our minimum population size estimate of *C. setosus* fluctuated over the 6 mark-recapture events. Our observed fluctuations were greater than the minimum population estimates; thus, our results suggest the population may be vulnerable to extirpation given catastrophic events. However, vulnerability is common when populations are isolated (Bland, 2017). Despite the general fragility of isolated populations, our minimum population size in 2010 overlapped with our estimate from 2019 suggesting some population stability that has been demonstrated in other cave-dwelling populations (e.g., Hobbs, 1978). Seasonal variation in the relative abundance of cave organisms is typical (Barr, 1967) and high numbers of individuals can be difficult to obtain (Cooper and Cooper, 1997; Miller and Niemiller, 2008), making longer term studies of cave species valuable. If future population estimates are desired for comparison to our estimates, studies would benefit from the increasing array of diminutive tags available for recapture studies over longer time periods (e.g., passive integrated transponders (Musselman et al., 2017), visible implant elastomer tags (Bolland et al., 2009; Venarsky et al., 2012), and p-Chips (Tenczar et al., 2014; Moore and Brewer, 2021)). Moreover, quantitative advancements that facilitate more robust mark-recapture designs would be beneficial to improving our understanding of population fluctuations (e.g., Royle-Nichols (Nakashima, 2020)). Lastly, estimating detection bias in population estimates would be desirable (Royle, 2004), but we were unable to do so because habitat was not measured on most of the repeat surveys.

Our results reflect the life history and basic ecology of a single population of *C. setosus* but adds to the growing body of literature necessary to conserve cave crayfishes and overcome our limited knowledge of species traits (Mammola et al., 2019). It is important to recognize that populations have genetic differences, which may translate into phenotypic or life history differences. Therefore, it is beneficial if future studies consider investigating cave crayfish population traits to assess generalizations that can be extended to other cave systems. We found that males can reproduce when their CLs reach 18 mm, and these data can be used in population models to predict changes when different management options are applied (e.g., Crouse et al., 1987). Conservation efforts focused on maintaining the natural habitat within




Figure 5. Photomicrograph of a bristly cave crayfish *Cambarus setosus* gastric mill section from a 30 mm carapace length individual collected from Smallin Civil War Cave, Christian County, Missouri, United States. No clear yearly bands are visible; however, faint bands can be seen and are denoted by the arrow. We hypothesize that these bands are sub-yearly in nature and may represent unidentified cycles such as feeding or temperature (Wright et al., 2002; Mouser et al., 2020).

the cave would be advantageous given the crayfish appear to use many different habitat components depending on their sex, the time of year, and location within the cave. Knowing the reproduction timing of crayfish can also aid in management decisions. For example, environmental DNA surveys would be most effective when they coincide with reproductive periods (e.g., de Souza et al., 2016). In contrast, it may be beneficial to avoid recreational caving during reproduction to avoid crushing crayfish which is a significant source of mortality (Graening et al., 2006).

ACKNOWLEDGEMENTS

This research was a contribution of the Oklahoma Cooperative Fish and Wildlife Research Unit (U.S. Geological Survey, Oklahoma Department of Wildlife Conservation, Oklahoma State University, and Wildlife Management Institute cooperating). Project funding was provided by the U.S. Fish and Wildlife Service (G15AC00021). Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. We would like to thank Kevin Bright for his continued support and enthusiasm for the project. Kevin provided access to Smallin Cave, assisted during several surveys, and shared photographs and anecdotal comments relevant to our understanding of the natural history of bristly cave crayfish *Cambarus setosus*. We would also like to acknowledge Jon Beard, members of the Springfield Plateau Grotto, Robert Mollenhauer, and students from Missouri Western State University, Reis Biological Station, and Oklahoma State University, who all provided technical assistance.

REFERENCES

- Adamski, J.C., 2000, Geochemistry of the Springfield Plateau aquifer of the Ozark Plateaus Province in Arkansas, Kansas, Missouri and Oklahoma, USA: Hydrological Processes, v. 14, p. 849–866. [https://doi.org/10.1002/\(SICI\)1099-1085\(20000415\)14:5<849::AID-HYP973>3.0.CO;2-7](https://doi.org/10.1002/(SICI)1099-1085(20000415)14:5<849::AID-HYP973>3.0.CO;2-7)
- Barr, T.C., Jr., 1967, Observations on the ecology of caves: The American Naturalist, v. 101, p. 475–491. <https://doi.org/10.1086/282512>
- Bell, G., 1974, Population estimates from recapture studies in which no recaptures have been made: Nature, v. 248, p. 616. <https://doi.org/10.1038/248616a0>
- Bland, L.M., 2017, Global correlates of extinction risk in freshwater crayfish: Animal Conservation, v. 20, p. 532–542. <https://doi.org/10.1111/acv.12350>
- Bolland, J.D., Cowx, I.G., and Lucas, M.C., 2009, Evaluation of VIE and PIT tagging methods for juvenile cyprinid fishes: Journal of Applied Ichthyology, v. 25, p. 381–386. <https://doi.org/10.1111/j.1439-0426.2009.01261.x>
- Boulton, A.J., Fenwick, G.D., Hancock, P.J., and Harvey, M.S., 2008, Biodiversity, functional roles and ecosystem services of groundwater invertebrates: Invertebrate Systematics v. 22, p. 103–116. <https://doi.org/10.1071/IS07024>
- Burnham, K.P., and Anderson, D.R., 2001, Kullback–Leibler information as a basis for strong inference in ecological studies: Wildlife Research, v. 28, p. 111–119. <https://doi.org/10.1071/WR99107>
- Caine, E., 1978, Comparative ecology of epigean and hypogean crayfish (Crustacea: Cambaridae) from northwestern Florida: The American Midland Naturalist, v. 99, p. 315–329. <https://doi.org/10.2307/2424809>
- Cooper, J.E., and Cooper, M.R., 1997, A new species of crayfish of the genus *Orconectes*, subgenus *Orconectes* (Decapoda: Cambaridae), endemic to Shelta Cave, Huntsville, Alabama: Journal of Cave and Karst Studies, v. 59, p. 119–127.
- Crandall, K.A., 2016, Collecting and processing freshwater crayfishes: Journal of Crustacean Biology, v. 36, p. 761–766. <https://doi.org/10.1163/1937240X-00002466>
- Crouse, D.T., Crowder, L.B., and Caswell H., 1987, A stage-based population model for loggerhead sea turtles and implications for conservation: Ecology, v. 68, p. 1412–1423. <https://doi.org/10.2307/1939225>
- Danielopol, D.L., and Griebler, C., 2008, Changing paradigms in groundwater ecology – from the 'living fossils' tradition to the 'new groundwater ecology': International Review of Hydrobiology, v. 93, p. 565–577. <https://doi.org/10.1002/iroh.200711045>
- de Souza, L.S., Godwin, J.C., Renshaw, M.A., and Larson, E., 2016, Environmental DNA (eDNA) detection probability is influenced by seasonal activity of organisms: PLoS ONE, v. 11, e0165273. <https://doi.org/10.1371/journal.pone.0165273>
- DiStefano, R.J., 2005, Trophic interactions between Missouri Ozarks stream crayfish communities and sport fish predators: increased abundance and size structure of predators cause little change in crayfish community densities: Missouri Department of Conservation Dingell-Johnson Project F-1-R-054, Study S-41, Job 4, Final Report.
- DiStefano, R.J., Black, T.R., Herleth-King, S.S., Kanno, Y., and Mattingly H.L., 2013, Life histories of two populations of the imperiled crayfish *Orconectes (Procericambarus) williamsi* (Decapoda: Cambaridae) in Southwestern Missouri, U.S.A: Journal of Crustacean Biology, v. 33, p. 15–24. <https://doi.org/10.1163/1937240X-00002109>
- DiStefano, R.J., Westhoff, J.T., Ames, C.W., and Rosenberger, A.E., 2016, Life history of the vulnerable endemic crayfish *Cambarus (Erebicambarus) maculatus* Hobbs and Pflieger, 1988 (Decapoda: Astacoidea: Cambaridae) in Missouri, USA: Journal of Crustacean Biology, v. 36, p. 615–627. <https://doi.org/10.1163/1937240X-00002472>
- DiStefano, R.J., Ashley, D., Brewer, S.K., Mouser, J.B., and Niemiller, M.L., 2020, Preliminary investigation of the critically imperiled Caney Mountain cave crayfish *Orconectes stygocaneyi* Hobbs III, 2001 (Decapoda: Cambaridae) in Missouri, USA: Freshwater Crayfish, v. 25, p. 47–57. <https://doi.org/10.5869/fc.2020.v25-1.047>
- DiStefano, R., Thoma, R.F., and Cordeiro, J., 2021, *Cambarus setosus* (amended version of 2010 assessment). The IUCN Red List of Threatened Species 2021: <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T153995A198269186.en>
- Dyer, J.J., Worthington, T.A., and Brewer, S.K., 2015, Response of crayfish to hyporheic water availability and excess sedimentation: Hydrobiologia, v. 747, p. 147–157. <https://doi.org/10.1007/s10750-014-2126-8>
- Dyer, J.J., Mouser, J., and Brewer S.K., 2016, Habitat use and growth of the western painted crayfish *Orconectes palmeri longimanus* (Faxon, 1898) (Decapoda: Cambaridae): Journal of Crustacean Biology, v. 36, p. 172–179. <https://doi.org/10.1163/1937240X-00002417>
- Edwards, A.W., 1974, Population estimates from recapture studies: Nature, v. 252, p. 509–510. <https://doi.org/10.1038/248616a0>
- Faraway, J.J., 2005, Extending the Linear Model with R: Generalized Linear, Mixed Effects and Nonparametric Regression Models: Boca Raton, CRC Press.
- Fero, K., and Moore, P.A., 2008, Social spacing of crayfish in natural habitats: what role does dominance play?: Behavioral Ecology and Sociobiology, v. 62, p. 1119–1125. <https://doi.org/10.1007/s00265-007-0540-x>

- Freund, R.J., and Wilson, W.J., 2003, *Statistical Methods* (second edition): San Diego, Academic Press.
- Friedenberg, N.A., Hoover, J.J., Boysen, K., and Killgore, K.J., 2018, Estimating abundance without recaptures of marked pallid sturgeon in the Mississippi River: *Conservation Biology*, v. 32, p. 457–465. <https://doi.org/10.1111/cobi.12972>
- Gardner, J.E., 1986, *Invertebrate Fauna from Missouri Caves and Springs*: Jefferson City Missouri Department of Conservation.
- Gelman, A., and Hill, J., 2007, *Data Analysis Using Regression and Multilevel/Hierarchical Models*: Cambridge, UK, Cambridge University Press.
- Graening, G.O., Horton, H.H., III, Slay, M.E., Elliott, W.R., and Brown, A.V., 2006, Status update for bristly cave crayfish, *Cambarus setosus* (Decapoda: Cambaridae), and range extension into Arkansas: *The Southwestern Naturalist*, v. 51, p. 382–392. [https://doi.org/10.1894/0038-4909\(2006\)51\[382:SUFBC\]2.0.CO;2](https://doi.org/10.1894/0038-4909(2006)51[382:SUFBC]2.0.CO;2)
- Griebler, C., Malard, F., and Lefebvre, T., 2014, Current developments in groundwater ecology from biodiversity to ecosystem function and services: *Current Opinion in Biotechnology*, v. 27, p. 159–167. <https://doi.org/10.1016/j.copbio.2014.01.018>
- Hobbs, H.H., III, 1978, Studies of the cave crayfish, *Orconectes inermis inermis* Cope (Decapoda, Cambaridae). Part IV: Mark-recapture procedures for estimating population size and movements of individuals: *International Journal of Speleology*, v. 10, p. 303–322. <http://dx.doi.org/10.5038/1827-806X.10.3.7>
- Jegla, T.C., 1966., Reproductive and molting cycles in cave crayfish: *Biological Bulletin*, v. 130, p. 345–358. <https://doi.org/10.2307/1539741>
- Jegla, T.C., and Poulson, T.L., 1970, Circannian rhythms—I. Reproduction in the cave crayfish, *Orconectes pellucidus inermis*: *Comparative Biochemistry and Physiology*, v. 33, p. 347–355. [https://doi.org/10.1016/0010-406X\(70\)90355-5](https://doi.org/10.1016/0010-406X(70)90355-5)
- Larson, E.R., and Olden, J.D., 2010, Latent extinction and invasion risk of crayfishes in the southeastern United States: *Conservation Biology*, v. 24, p. 1099–1110. <https://doi.org/10.1111/j.1523-1739.2010.01462.x>
- Leland, J.C., Bucher, D.J., and Coughran, J., 2015, Direct age determination of a subtropical freshwater crayfish (Redclaw, *Cherax quadricarinatus*) using ossicular growth marks: *PLoS ONE*, v. 10, e0134966. <https://doi.org/10.1371/journal.pone.0134966>
- Light, T., 2003, Success and failure in a lotic crayfish invasion: the roles of hydrologic variability and habitat alteration: *Freshwater Biology*, v. 48, p. 1886–1897. <https://doi.org/10.1046/j.1365-2427.2003.01122.x>
- Mammola, S., et al., 2019, Scientists' warning on the conservation of subterranean ecosystems: *BioScience*, v. 69, p. 641–650. <https://doi.org/10.1093/biosci/biz064>
- Marquart, D., 1979, *The troglotic crayfish of Missouri [M.S. thesis]*: Warrensburg, Central Missouri State University.
- Miller, B.T., and Niemiller, M.L., 2008, Distribution and relative abundance of Tennessee cave salamanders (*Gyrinophilus pallescens* and *Gyrinophilus gulolineatus*) with an emphasis on Tennessee populations: *Herpetological Conservation and Biology*, v. 3, p. 1–20.
- Missouri Department of Conservation, 2021, *Missouri Species and Communities of Conservation Concern Checklist*: Jefferson City Missouri, Missouri Department of Conservation.
- Momot, W.T., 1995, Redefining the role of crayfish in aquatic ecosystems: *Reviews in Fisheries Science*, v. 3, p. 33–63. <https://doi.org/10.1080/10641269509388566>
- Moore, D., and Brewer, S.K., 2021, Evaluation of VIE, PIT, and p-Chip tagging methods in small bodied minnow species: *North American Journal of Fisheries Management*, v. 41, p. 1066–1078. <https://doi.org/10.1002/nafm.10607>
- Moore, M.J., DiStefano, R.J., and Larson, E.R., 2013, An assessment of life-history studies for USA and Canadian crayfishes: Identifying biases and knowledge gaps to improve conservation and management: *Freshwater Science*, v. 32, p. 1276–1287. <https://doi.org/10.1899/12-158.1>
- Mouser, J.M., Ashley, D.C., Aley, T., and Brewer, S.K., 2019, Subterranean invasion by gapped ringed crayfish: Effectiveness of a removal effort and barrier installation: *Diversity*, v. 11, 3. <https://doi.org/10.3390/d11010003>
- Mouser, J.M., Glover, J., and Brewer, S.K., 2020, Gastric mill age estimates for North American crayfish and the influence of temperature: *Freshwater Crayfish*, v. 25, p. 59–67. <https://doi.org/10.5869/fc.2020.v25-1.059>
- Musselman, W.C., Worthington, T.A., Mouser, J., Williams, D.M., and Brewer S.K., 2017, Passive integrated transponder tags: review of studies on warmwater fishes with notes on additional species: *Journal of Fish and Wildlife Management*, v. 8, p. 353–364. <https://doi.org/10.3996/122016-JFWM-091>
- Nakashima, Y., 2020, Potentiality and limitations of N-mixture and Royle-Nichols models to estimate animal abundance based on noninstantaneous point surveys: *Population Ecology*, v. 62, p. 151–157. <https://doi.org/10.1002/1438-390X.12028>
- NatureServe, 2009, Bristly cave crayfish *Cambarus setosus*: https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.116048/Cambarus_setosus (visited July 2022)
- Paine, R.T., 1969, A note on trophic complexity and community stability: *The American Naturalist*, v. 103, p. 91–93. <https://doi.org/10.1086/282586>
- Panfili, J., de Pontual, H., Troadec, H., and Wright, P.J., 2002. *Manual of Fish Sclerochronology*: Best, France, Ifremer.
- Pflieger, W.L., 1996, *The Crayfishes of Missouri*: Jefferson City, Missouri Department of Conservation.
- Poulson, T.L., 2012, Cave ecosystems, in White, W.B., and Culver, D.C., eds., *Encyclopedia of Caves* (second edition): Cambridge, Academic Press, p. 323–334.
- R Core Team, 2020, *R: A language and environment for statistical computing*: Foundation for Statistical Computing: <https://cran.r-project.org/doc/manuals/r-release/fullrefman.pdf> (accessed July 2022).
- Ramalho, R.O., McClain, W.R., and Anastácio, P.M., 2010, An effective and simple method of temporarily marking crayfish: *Freshwater Crayfish*, v. 17, p. 57–60. <https://doi.org/10.5869/fc.2010.v17.57>
- Renai, B., Bertocchi, S., Brusconi, S., Gherardi, F., Grandjean, F., Lebboroni, M., Parinet, B., Grosset, C.S., and Trouilhe, M.C., 2006, Ecological characterisation of streams in Tuscany (Italy) for the management of the threatened crayfish *Austropotamobius pallipes* complex: *Bulletin Français de la Pêche et de la Pisciculture*, v. 380381, p. 1095–1114. <https://doi.org/10.1051/kmae:2006013>
- Royle, J.A., 2004, N-mixture models for estimating population size from spatially replicated counts: *Biometrics*, v. 60, p. 108–115. <https://doi.org/10.1111/j.0006-341X.2004.00142.x>
- Schneider, R.A.Z., Schneider, R.W.S., and Moore, P.A., 1999, Recognition of dominance status by chemoreception in the red swamp crayfish, *Procambarus clarkia*: *Journal of Chemical Ecology*, v. 25, p. 781–794. <https://doi.org/10.1023/A:1020888532513>
- Simon, K.S., 2012, Cave ecosystems, in White, W.B., and Culver, D.C., eds., *Encyclopedia of Caves* (second edition): Cambridge, Academic Press, p. 99–102.
- Smith, G.R.T., Learner, M.A., Slater, F.M., and Foster, J., 1996, Habitat features important for the conservation of the native crayfish *Austropotamobius pallipes* in Britain: *Biological Conservation*, v. 75, p. 239–246. [https://doi.org/10.1016/0006-3207\(95\)00073-9](https://doi.org/10.1016/0006-3207(95)00073-9)
- Streissl, F., and Hödl, W., 2002, Habitat and shelter requirements of the stone crayfish, *Austropotamobius torrentium* Schrank: *Hydrobiologia*, v. 477, p. 195–199. <https://doi.org/10.1023/A:1021094309738>

- Taylor, C.A., Schuster, G.A., Cooper, J.E., DiStefano, R.J., Eversole, A.G., Hamr, P., Hobbs, H.H., III, Robinson, H.W., Skelton, C.E., and Thoma, R.F., 2007, A reassessment of the conservation status of crayfishes in the United States and Canada after 10+ years of increased awareness: *Fisheries*, v. 32, p. 372–389. [https://doi.org/10.1577/1548-8446\(2007\)32\[372:AROTCS\]2.0.CO;2](https://doi.org/10.1577/1548-8446(2007)32[372:AROTCS]2.0.CO;2)
- Taylor, C.A., DiStefano, R.J., Larson, E.R., and Stoeckel, J., 2019, Towards a cohesive strategy for the conservation of the United States' diverse and highly endemic crayfish fauna: *Hydrobiologia*, v. 846, p. 39–58. <https://doi.org/10.1007/s10750-019-04066-3>
- Tenczar, P., Lutz, C.C., Rao, V.D., Goldenfeld, N., and Robinson, G.E., 2014, Automated monitoring reveals extreme interindividual variation and plasticity in honeybee foraging activity levels: *Animal Behaviour*, v. 95, p 41–48. <https://doi.org/10.1016/j.anbehav.2014.06.006>
- Unklesbay, A.G, and Vineyard, J.D. 1992. *Missouri Geology: Three Billion Years of Volcanoes, Seas, Sediments, and Erosion*: Missouri, University of Missouri Press.
- Venables, W.N., and Ripley, B.D., 2002, *Modern Applied Statistics with S* (fourth edition): New York, Springer.
- Venarsky, M.P., Huryn, A.D., and Benstead, J.P., 2012, Re-examining extreme longevity of the cave crayfish *Orconectes australis* using new mark–recapture data: A lesson on the limitations of iterative size-at-age models: *Freshwater Biology*, v. 57, p. 1471–1481. <https://doi.org/10.1111/j.1365-2427.2012.02812.x>
- Woods, A.J., Omernik, J.M., Butler, D.R., Ford, J.G., Henley, J.E., Hoagland, B.W., Arndt, D.S., and Moran B.C., 2005, Ecoregions of Oklahoma (color poster with map, descriptive text, summary tables, and photographs): Virginia, U.S. Geological Survey.
- Wright, P.J., Panfili, J., Morales-Nin, B., and Geffen, A.J, 2002, Types of calcified structures: Otoliths, *in* Panfili J. de Pontual H., Troadec H., and Wright, P.J. eds., *Manual of Fish Sclerochronology*: Best, Ifremer, p. 31–56.
- Yau, C., 2013, Multinomial goodness of fit: <http://www.r-tutor.com/elementary-statistics/goodness-fit/multinomial-goodness-fit> (accessed October 2020).
- Zar, J.H., 1999, *Biostatistical Analysis* (fourth edition): Upper Saddle River, Prentice Hall.